

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



1215-1171
2513-1158

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07H 21/04, C12Q 1/68 A61K 48/00	A1	(11) International Publication Number: WO 94/06815 (43) International Publication Date: 31 March 1994 (31.03.94)
(21) International Application Number: PCT/US93/08367 (22) International Filing Date: 3 September 1993 (03.09.93) (30) Priority data: 07/943,516 11 September 1992 (11.09.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/943,516 (CIP) Filed on 11 September 1992 (11.09.92) (71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]: 2280 Faraday Avenue, Carlsbad, CA 92008 (US).	(72) Inventors; and (75) Inventors/Applicants (for US only): MANOHARAN, Mu- thiah [IN/US]: 7634 Reposado Drive, Carlsbad, CA 92009 (US). COOK, P. Dan [US/US]: 7340 Bolero Street, Carlsbad, CA 92009 (US). (74) Agents: CALDWELL, John. W. et al.; Woodcock Wash- burn Kurtz Mackiewicz & Norris, One Liberty Place - 46th Floor, Philadelphia, PA 19103 (US). (81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN. European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: OLIGONUCLEOTIDE AND NUCLEOTIDE AMINE ANALOGS. METHODS OF SYNTHESIS AND USE (57) Abstract Oligonucleotide and nucleotide amine analogs and methods of preparing and using these compounds are provided by the present invention.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TC	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

- 1 -

OLIGONUCLEOTIDE AND NUCLEOTIDE AMINE ANALOGS, METHODS OF
SYNTHESIS AND USE

FIELD OF THE INVENTION

This invention relates to novel amine-containing compounds useful for therapeutics and methods of making and
5 using the same.

BACKGROUND OF THE INVENTION

It is well known that most of the bodily states in mammals including most disease states, are effected by proteins. Such proteins, either acting directly or through
10 their enzymatic functions, contribute in major proportion to many diseases in animals and man. Classical therapeutics has generally focused upon interactions with such proteins in efforts to moderate their disease causing or disease potentiating functions. Recently, however, attempts have been
15 made to moderate the actual production of such proteins by interactions with molecules that direct their synthesis, intracellular RNA. These interactions involved the binding of complementary "antisense" oligonucleotides or their analogs to the transcellular RNA in a sequence specific fashion such as by
20 Watson-Crick base pairing interactions.

The pharmacological activity of antisense compounds, as well as other therapeutics, depends on a number of factors that influence the effective concentration of these agents at specific intracellular targets. One important factor is the
25 ability of antisense compounds to traverse the plasma membrane of specific cells involved in the disease process.

Cellular membranes consist of lipid protein bilayers that are freely permeable to small, nonionic, lipophilic

- 2 -

compounds and inherently impermeable to most natural metabolites and therapeutic agents. Wilson, *Ann. Rev. Biochem.* 1978, 47, 933. The biological and antiviral effects of natural and modified oligonucleotides in cultured mammalian cells have been well documented, so it appears that these agents can penetrate membranes to reach their intracellular targets. Uptake of antisense compounds into a variety of mammalian cells, including HL-60, Syrian Hamster fibroblast, U937, L929, CV-1, and ATH8 cells has been studied using natural oligonucleotides and nuclease resistant analogs, such as alkyl triesters, Miller, et al., *Biochemistry* 1977, 16, 1988; methylphosphonates, Marcus-Sekura, et al., *Nuc. Acids Res.* 1987, 15, 5749 and Miller, et al., *Biochemistry* 1981, 20, 1874; and phosphorothioates, Ceruzzi, et al., *Nucleosides & Nucleotides* 1989, 8, 815; Miller, et al., *Biochemistry* 1987, 16, 1988; and Loke, et al., *Curr. Top. Microbiol. Immunol.* 1988, 141, 282.

Enhanced cellular uptake has previously been achieved by attachment of functional groups to the 3' and 5' end of oligonucleotides to enhance cellular uptake in specific cell types. Previous studies have shown that plasmid DNA complexed with an (asialo)glycoprotein-poly(L-lysine) conjugate, could be targeted to hepatocytes, which contain unique cell surface receptors for galactose-terminal (asialo)glycoproteins. Wu, et al., *Biochemistry* 1988, 27, 887. Other groups have synthesized oligodeoxyribonucleotides that have a 5'-attached alkylating agent and a 3' attached cholesterol moiety and determined that these modified oligonucleotides were taken up into cells more efficiently than control compounds without the steroid moiety. Zon, G. in *Oligodeoxynucleotides: Antisense Inhibitors of Gene Expression* 234-247, ed. J.S. Cohen (CRC Press, Boca Raton FL, 1989). Letsinger, et al., *Proc. Natl. Acad. Sci. U.S.A.* 1989, 86, 653, have also synthesized cholesteryl-conjugated phosphorothioates whose anti-HIV activity is significantly greater than natural oligonucleotides with the same sequence. Additional modifications include conjugation of

- 3 -

oligonucleotides to poly(L-lysine) alone. Stevenson, et al., *J. Gen. Virol* 1989, 70, 2673 and Lemaitre, et al., *Proc. Natl. Acad. Sci. U.S.A.* 1987, 84, 648. This modification enhanced the antiviral activity of the compound studied presumably due to increased cellular uptake imparted by the polycationic poly(L-lysine).

The conjugation of polyamines to oligonucleotides have been found to enhance cellular uptake of oligonucleotides, increased lipophilicity, cause greater cellular retention and increased distribution of the compound. Vasseur, *Nucleosides and Nucleotides* 1991, 10, 107 prepared abasic sites at different sites of oligothymidylates by acid hydrolysis. Thereafter the abasic sites were functionalized with functionalities such as 3-amino carbazole, 9-amino elipticine and psoralen. Vasseur, et al., also refer to unpublished results in which the functionalities spermidine and proflavin were employed. The abasic site was generated by one of the following three methods: (i) selective depurination by acid treatment in a pyrimidine-rich oligonucleotide having one purine in a chosen site, (ii) incorporating 2',3'-dideoxynebularine at the 5'-end with the nebularine phosphoramidite at the last step of the oligonucleotide synthesis, and subsequent acid treatment (30mM HCl at 37°C) to create an abasic site at 5' end (in this case the open-chain structure is $\text{CHO}-(\text{CH}_2)_2-\text{CHOH}-\text{CH}_2\text{O}$ -at the 5' end and the conjugate from the amine RNH_2 is $\text{RNH}-(\text{CH}_2)_3-\text{CHOH}-\text{CH}_2\text{O}$ -Oligo), and (iii) incorporating a protected abasic 2'-deoxy-D-ribofuranose nucleotide synthon that has a photo-labile O-nitrobenzyl group as the anomeric hydroxyl-protecting group in oligonucleotide synthesis and removing it prior to conjugation.

Groebke and Leumann used a silyl-protecting group at the anomeric center to generate the abasic site. 2'-Deoxy-5-O-dimethoxytrityl-D-ribofuranose was silylated at the 1-O-position using TBDMSCl and the silyl group was removed later by hydrolysis at pH 2.0 to yield the abasic site. Unfortunately, fluoride-ion-mediated deprotection of the silyl group caused a β -elimination and DNA degradation.

- 4 -

McLaughlin's group has utilized 1-(β -D-2-deoxyribo-
syl)-2-pyrimidone-based phosphoramidite to generate
abasic sites at pH 3.0. The N-glycosyl cleavage occurred,
however, slower in oligonucleotides than in parent nucleosides;
5 nearly 60 hours of acid treatment was necessary to generate 90%
abasic site formation. However, conjugation chemistry via
enzymatically generated abasic sites are unknown in the
literature.

Le Doan, et al., *Nucleic Acids Research* 1987, 15, 8643
10 teaches oligothymidylates covalently linked to porphyrins at
their 3' end via one of the linkers $-O-CH_2-CO-NH-(CH_2)_2-NH$ or
 $PO_4-(CH_2)_6-NH-$. Le Doan, et al., also used the linker PO_4-
 $(CH_2)_6-NH-$ to link porphyrins to the 5' end of
oligothymidylates. Another group, Summerton, et al., U.S.
15 Patent No. 5,034,506 issued July 23, 1991 teaches morpholino
subunits, linked together by uncharged, achiral linkages such
as amides. As described in PCT/US91/04086 filed June 10, 1991,
polyamines have also been linked at the 5' end of an
oligonucleotide at the 5' site of the sugar moiety of the
20 terminal nucleoside and at the 2-position carbon of the
heterocyclic base of 2'-deoxyadenosine, 2'-deoxyguanosines and
other purines and purine analogs by known procedures as
described in PCT/US/91/00243 filed January 11, 1991.

Novel amines and methods of preparing the same are
25 greatly needed in order to enhance cellular uptake of
oligonucleotides, increase lipophilicity, cause greater
cellular retention and increase distribution of the compound
within the cell. The present invention fulfills this need.

OBJECTS OF THE INVENTION

30 It is one object of the present invention to provide
novel amine-containing compounds useful in therapeutics.

It is a further object of the present invention to
provide methods of producing said novel compounds.

It is another object of the present invention to
35 provide methods of modulating the production of a protein by an
organism.

- 5 -

It is still a further object of the present invention to provide methods of treating a mammal suffering from a disease characterized by the undesired production of a protein.


It is yet a further object of the present invention to provide methods of diagnosing the presence of an RNA in a biological sample.

These and other objects will become apparent from the following description and accompanying claims.

SUMMARY OF THE INVENTION

The present invention provides compounds which may have enhanced efficacy as an antisense-based therapy. Compounds of the present invention can have enhanced cellular uptake, increased lipophilicity, cause greater cellular retention and demonstrate increased distribution. Furthermore the present invention provides simple methods for synthesis of these novel compounds.

In accordance with some embodiments of the present invention, compounds having the structure:




I

wherein R_1 and R_2 are independently H, a nucleotide, oligonucleotide, or an amine-containing species, and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic amine-containing species, and X is H, O- R_{11} , S- R_{11} , F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or

- 6 -

substituted lower alkyl, C₂ to C₁₀ straight or branched chain lower alkenyl or substituted lower alkenyl, C₃ to C₁₀ straight or branched chain lower alkynyl or substituted lower alkynyl, a ¹⁴C containing lower alkyl, lower alkenyl or lower alkynyl, a ¹⁴C containing C₇ to C₁₄ alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide, are provided.


In accordance with still other embodiments of the present invention, compounds having the structure:



II

wherein R₄ is an oligonucleotide and M is a pendent group having an amine-containing species attached thereto are provided.

Methods of preparing such compounds utilizing enzymatic reagents are also provided in some aspects of the invention. Thus compounds of Formula I may be prepared by methods comprising the steps of providing a synthon having the structure:




wherein R₁ and R₂ are independently H, a nucleotide, oligonucleotide or amine-containing species, and at least one of R₁ and R₂ is a purine containing oligonucleotide, and X is H, O-R₁₁, S-R₁₁, F, Cl, Br, CN, CF₃, OCF₃, OCN, SOCH₃, SO₂CH₃,

- 7 -

ONO₂, N₃, HN₂, heterocylcoalkyl, heterocycloalkaryl,
 aminoalkylamino, polyalkylamino, substituted silyl, a reporter
 molecule, an RNA cleaving group, a group for improving the
 pharmacokinetic properties of an oligonucleotide, or a group
 5 for improving the pharmacodynamic properties of an
 oligonucleotide wherein R₁₁ is H, C₁ to C₁₀ straight or branched
 chain lower alkyl or substituted lower alkyl, C₂ to C₁₀ straight
 or branched chain lower alkenyl or substituted lower alkenyl,
 C₃ to C₁₀ straight or branched chain lower alkynyl or
 10 substituted lower alkynyl, a ¹⁴C containing lower alkyl, lower
 alkenyl or lower alkynyl, C₇ to C₁₄ substituted or unsubstituted
 alkyaryl or aralkyl, a ¹⁴C containing C₇ to C₁₄ alkaryl or
 aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA
 cleaving group, a group for improving the pharmacokinetic
 15 properties of an oligonucleotide or a group for improving the
 pharmacodynamic properties of an oligonucleotide. Thereafter
 the synthon is reacted with R₃, wherein R₃ is a linear or cyclic
 amine-containing species, under reducing conditions to yield
 the final product.

20 Compounds of Formula II may also be prepared
 enzymatically by providing a starting material having the
 structure:




wherein R₄ is an oligonucleotide, R₁₂ is an oligonucleotide and
 B is urea or a heterocyclic base having a corresponding
 25 glycosylase and reacting the starting material with an
 endonuclease to generate a conjugated α,β-unsaturated system in
 the sugar residue of the 3' terminal nucleotide. Thereafter
 the compound having a conjugated α,β-unsaturated system is
 reacted with a pendent group containing a nucleophile
 30 functionality thereon. Following addition of the pendent group
 the double bond of the α,β system is reduced with a reducing
 agent. An amine-containing species may then be attached to the

- 8 -

pendent group via an alkylation reaction. Alternatively, an amine-containing species may be attached to a pendent group which is a bifunctional linker.


In accordance with still other embodiments of the present invention compounds having the structure:



III

wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic amine-containing species containing at least one non-amide nitrogen atom, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle, are provided. Methods of preparing compounds of Formula III are also provided in some aspects of the present invention comprising the steps of reacting an oligonucleotide having a 3' ribofuranosyl sugar with an oxidizing agent to produce an activated dialdehyde-terminated oligonucleotide and reacting said activated oligonucleotide with a linear or cyclic amine-containing species under reducing conditions to yield said compound.

In accordance with other aspects of the invention compounds having the structure:



IV

wherein B is a purine or pyrimidine heterocyclic base, R_8 and R_9 are independently H, PO_2^- , a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_8 and R_9 is a purine containing oligonucleotide, and at least one of R_8 , R_9 and A is a species comprising the formula L_1 - L_2 -polyamine wherein L_1 is an amino linker and L_2 is a heterobifunctional linker; and wherein if R_8 is not a purine containing

- 9 -

oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2 ; if R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH are provided.

5 Therapeutic and diagnostic methods are also encompassed by the present invention. Methods of modulating the production of protein by an organism comprising contacting an organism with a compound having the structure of Formula I, Formula II, Formula III or Formula IV are encompassed by some
10 embodiments of the present invention. In other aspects of the invention, methods of treating an animal having a disease characterized by undesired production of protein comprising contacting an animal with a compound having the structure of Formula I, Formula II, Formula III, or Formula IV in a
15 pharmaceutically acceptable carrier are provided. Still other methods of the present invention provide methods for detecting the presence or absence of an RNA in a biological sample suspected of containing said RNA are provided comprising contacting a sample with a compound having the structure of
20 Formula I, Formula II, Formula III or Formula IV wherein the compound is specifically hybridizable with the RNA and detecting the presence or absence of hybridization of the compound to the sample wherein hybridization is indicative of the presence of RNA in the sample.

25 BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a schematic representation of one preferred syntheses of compounds of Formula I.


Figure 2 is a schematic representation of one preferred syntheses of compounds of Formula II.

30 Figure 3 is a schematic representation of one preferred syntheses of compounds of Formula III.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention provides novel amine compounds useful for antisense therapy. In one embodiment of the present
35 invention compounds having the structure:

- 10 -



I

wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide, or an amine-containing species, and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic amine-containing species, and X is H, O- R_{11} , S- R_{11} , F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkyaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide, are provided. In some embodiments of the present invention both R_1 and R_2 are oligonucleotides, at least one of which includes at least one purine nucleotide.

In the context of this invention, the term "oligonucleotide" refers to a polynucleotide formed from naturally occurring bases, such as purine and pyrimidine heterocycles, and furanosyl groups joined by native phosphodiester bonds. This term effectively refers to naturally occurring species or synthetic species formed from

- 11 -

naturally occurring subunits or their close homologs. The term "oligonucleotide" may also refer to moieties which have portions similar to naturally occurring oligonucleotides but which have non-naturally occurring portions. Thus, 5 oligonucleotides may have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur-containing species which are known for use in the art. In accordance with some preferred embodiments, at least some of the phosphodiester bonds of the oligonucleotide have 10 been substituted with a structure which functions to enhance the stability of the oligonucleotide or the ability of the oligonucleotide to penetrate into the region of cells where the viral RNA is located. It is preferred that such substitutions comprise phosphorothioate bonds, phosphotriesters, methyl 15 phosphonate bonds, short chain alkyl or cycloalkyl structures or short chain heteroatomic or heterocyclic structures. Most preferred are $\text{CH}_2\text{-NH-O-CH}_2$, $\text{CH}_2\text{-N(CH}_3\text{)-O-CH}_2$, $\text{CH}_2\text{-O-N(CH}_3\text{)-CH}_2$, $\text{CH}_2\text{-N(CH}_3\text{)-N(CH}_3\text{)-CH}_2$ and $\text{O-N(CH}_3\text{)-CH}_2\text{-CH}_2$ structures where phosphodiester is O-P-O-CH_2). Also preferred are morpholino 20 structures. Summerton, et al., U.S. 5,034,506 issued July 23, 1991. In other preferred embodiments, such as the protein-nucleic acid (PNA) backbone, the phosphodiester backbone of the oligonucleotide may be replaced with a polyamide backbone, the bases being bound directly or indirectly to the aza nitrogen 25 atoms of the polyamide backbone. see, e.g., Nielsen, et al., Science 1991, 254 1497 and WO 92/20702, published November 26, 1992. In accordance with other preferred embodiments, the phosphodiester bonds are substituted with other structures which are, at once, substantially non-ionic and non-chiral, or 30 with structures which are chiral and enantiomerically specific. Still other linkages include the those disclosed in United States Patent Applications Serial Number 566,836, filed August 13, 1990, entitled Novel Nucleoside Analogs; Serial Number 703,619, filed May 21, 1991, entitled Backbone Modified 35 Oligonucleotide Analogs; Serial Number 903,160, filed June 24, 1992, entitled Heteroatomic Oligonucleoside Linkages; Serial Number PCT/US92/04294, filed May 21, 1992, entitled Backbone

- 12 -

Modified Oligonucleotides; and Serial Number PCT/US92/04305, all assigned to the assignee of this invention. Persons of ordinary skill in the art will be able to select other linkages for use in practice of the invention.

5 Oligonucleotides may also include species which include at least some modified base forms. Thus, purines and pyrimidines other than those normally found in nature may be so employed. For example, deaza or aza purines and pyrimidines may be used in place of naturally purine or pyrimidine bases
10 and pyrimidine bases having substituent groups at the 5- or 6-positions; purine bases having altered or replacement substituent groups at the 2-, 6- or 8- positions are also provided in some aspects of the present invention. Similarly, modifications on the furanosyl portion of the nucleotide
15 subunits may also be effected, as long as the essential tenets of this invention are adhered to. Examples of such modifications are 2'-O-alkyl- and 2'-halogen-substituted nucleotides. Some specific examples of modifications at the 2' position of sugar moieties which are useful in the present
20 invention are OH, SH, SCH₃, F, OCN, O(CH₂)_nNH₂, O(CH₂)_nCH₃ where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl or aralkyl; Cl, Br, CN, CF₃, OCF₃, O-, S-, or N- alkyl; O-, S-, or N-alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino;
25 polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar
30 properties. Sugar mimetics such as cyclobutyls may also be used in place of the pentofuranosyl group. Exemplary modifications are disclosed in United States Patent Applications: Serial Number 463,358, filed January 11, 1990, entitled Compositions And Methods For Detecting And Modulating RNA
35 Activity; Serial Number 566,977, filed August 13, 1990, entitled Sugar Modified Oligonucleotides That Detect And Modulate Gene Expression; Serial Number 558,663, filed July 27,

- 13 -

1990, entitled Novel Polyamine Conjugated Oligonucleotides; Serial Number 558,806, filed July 27, 1991, entitled Nuclease Resistant Pyrimidine Modified Oligonucleotides That Detect And Modulate Gene Expression; and Serial Number PCT/US91/00243, 5 filed January 11, 1991, entitled Compositions and Methods For Detecting And Modulating RNA Activity; Serial Number 777,670, filed October 15, 1991, entitled Oligonucleotides Having Chiral Phosphorus Linkages; Serial Number 814,961, filed December 24, 1991, entitled Gapped 2' Modified Phosphorothioate 10 Oligonucleotides; Serial Number 808,201, filed December 13, 1991, entitled Cyclobutyl Oligonucleotide Analogs; and Serial Number 782,374, filed 782,374, entitled Derivatized Oligonucleotides Having Improved Uptake & Other Properties, all assigned to the assignee of this invention. The disclosures of 15 all of the above noted patent applications are incorporated herein by reference. Oligonucleotides may also comprise other modifications consistent with the spirit of this invention. Such oligonucleotides are best described as being functionally interchangeable with yet structurally distinct from natural 20 oligonucleotides. All such oligonucleotides are comprehended by this invention so long as they effectively function as subunits in the oligonucleotide. Thus, purine containing oligonucleotide are oligonucleotides comprising at least one purine base or analog thereof. In other embodiments of the 25 present invention compounds of the present invention may be "subunits" of a species comprising two or more compounds of the present invention which together form a single oligonucleotide.

Oligonucleotides of the present invention may be naturally occurring or synthetically produced and may range in 30 length from about 8 to about 50 nucleotides. In more preferred embodiments of the present invention said oligonucleotides may be from 8 to 40 nucleotides in length. Most preferably, oligonucleotides of the present invention may be from 12 to about 20 nucleotides in length.

35 Amine-containing species according to the invention are aromatic species containing a single nitrogen atom or non-aromatic species containing one or more nitrogen atoms (i.e.,

- 14 -

polyamine species). Amine-containing species can be linear (including straight-chain and branched) or cyclic. Cyclic amine-containing species can be aromatic or non-aromatic. Representative amine-containing species include amino acids, polypeptides, hydrazide salts of organic acids, including one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, cross-linking agents, peptide nucleic acids (PNA) and PEG (polyethylene glycol containing)-amines attached to at least one of the nitrogen atoms of said amine-containing species.

Polyamine species according to the invention are those that contain a plurality of nitrogen atoms. Polyamines include primary amines, hydrazines, semicarbazines, thiosemicarbazines and similar nitrogenous species. Such species can be symmetrical species such as polyamine-containing polymers or they can be unsymmetrical wherein the amine functionalities of the polyamine are separated in space by different moieties. In addition to carbon atoms other atomic species such as nitrogen and sulfur may also be incorporated into the polyamine species. In some preferred embodiments of the invention, at least one nitrogen atom of the polyamine has a free electron pair.

Preferred as polyamine species are species that range in length from about 3 to about 20 units. More preferably species having at least one nitrogen atom have the general formula $H_2N[(CH_2)_nNH]_m$ wherein n is an integer between 2 and 8 and m is an integer between 1 and 10. These species can be linear or cyclic. Cyclic amines would include crown amines ("cyclams") and mixed crown amines/crown ethers.

Other suitable amine-containing species according to the invention include C_1 - C_{20} straight chain alkylamine, C_1 - C_{20} straight chain substituted alkylamine, C_2 - C_{50} branched chain alkylamine, C_2 - C_{50} branched chain substituted alkylamine, C_3 - C_{50} cyclic alkylamine, C_3 - C_{50} cyclic substituted alkylamine, C_2 - C_{20} straight chain alkenylamine, C_2 - C_{20} straight chain substituted alkenylamine, C_3 - C_{50} branched chain alkenylamine, C_3 - C_{50} branched chain substituted alkenylamine, C_3 - C_{50} cyclic alkenylamine, C_3 -

- 15 -

- C₅₀ cyclic substituted alkenylamine, C₂-C₂₀ straight chain alkynylamine, C₂-C₂₀ straight chain substituted alkynylamine, C₃-C₅₀ branched chain alkynylamine, C₃-C₅₀ branched chain substituted alkynylamine, C₃-C₅₀ cyclic alkynylamine, C₃-C₅₀
- 5 cyclic substituted alkynylamine, C₁-C₂₀ straight chain alkylhydrazine, C₁-C₅₀ straight chain substituted alkylhydrazine, C₂-C₅₀ branched chain alkylhydrazine, C₂-C₅₀ branched chain substituted alkylhydrazine, C₃-C₅₀ cyclic hydrazoalkane, C₃-C₅₀ cyclic substituted hydrazoalkane, C₂-C₂₀
- 10 straight chain alkenylhydrazine, C₂-C₂₀ straight chain substituted alkenylhydrazine, C₃-C₅₀ branched chain alkenylhydrazine, C₃-C₅₀ branched chain substituted alkenylhydrazine, C₃-C₅₀ cyclic hydrazoalkene, C₃-C₅₀ cyclic substituted hydrazoalkene, C₂-C₂₀ straight chain
- 15 alkynylhydrazine, C₂-C₂₀ straight chain substituted alkynylhydrazine, C₃-C₅₀ branched chain alkynylhydrazine, C₃-C₅₀ branched chain substituted alkynylhydrazine, C₃-C₅₀ cyclic hydrazoalkyne, C₃-C₅₀ cyclic substituted hydrazoalkyne, C₁-C₂₀ straight chain alkylhydroxyamine, C₁-C₂₀ straight chain
- 20 substituted alkylhydroxyamine, C₂-C₅₀ branched chain alkylhydroxyamine, C₂-C₅₀ branched chain substituted alkylhydroxyamine, C₃-C₅₀ cyclic oxyalkylamine, C₃-C₅₀ cyclic substituted oxyalkylamine, C₂-C₂₀ straight chain alkenylhydroxyamine, C₂-C₂₀ straight chain substituted
- 25 alkenylhydroxyamine, C₃-C₅₀ branched chain alkenylhydroxyamine, C₃-C₅₀ branched chain substituted alkenylhydroxyamine, C₃-C₅₀ cyclic oxyalkenylamine, C₃-C₅₀ cyclic substituted oxyalkenylamine, C₂-C₂₀ straight chain alkynylhydroxyamine, C₂-C₂₀ straight chain substituted alkynylhydroxyamine, C₃-C₅₀
- 30 branched chain alkynylhydroxyamine, C₃-C₅₀ branched chain substituted alkynylhydroxyamine, C₃-C₅₀ cyclic oxyalkynylamine, C₃-C₅₀ cyclic substituted oxyalkynylamine, C₁-C₂₀ straight chain alkylsemicarbazide, C₁-C₂₀ straight chain substituted alkylsemicarbazide, C₂-C₅₀ branched chain alkylsemicarbazide,
- 35 C₂-C₅₀ branched chain substituted alkylsemicarbazide, C₃-C₅₀ cyclic alkylsemicarbazide, C₃-C₅₀ cyclic substituted alkylsemicarbazide, C₂-C₂₀ straight chain alkenylsemicarbazide,

- 16 -


- C_2-C_{20} straight chain substituted alkenylsemicarbazide, C_3-C_{50} branched chain alkenylsemicarbazide, C_3-C_{50} branched chain substituted alkenylsemicarbazide, C_3-C_{50} cyclic alkenylsemicarbazide, C_3-C_{50} cyclic substituted alkenylsemicarbazide, C_2-C_{20} straight chain alkynylsemicarbazide, C_2-C_{20} straight chain substituted alkynylsemicarbazide, C_3-C_{50} branched chain alkynylsemicarbazide, C_3-C_{50} branched chain substituted alkynylsemicarbazide, C_3-C_{50} cyclic alkynylsemicarbazide, C_3-C_{50} cyclic substituted alkynylsemicarbazide, C_1-C_{20} straight chain alkylthiosemicarbazide, C_1-C_{20} straight chain substituted alkylthiosemicarbazide, C_2-C_{50} branched chain alkylthiosemicarbazide, C_2-C_{50} branched chain substituted alkylthiosemicarbazide, C_3-C_{50} cyclic alkylthiosemicarbazide, C_3-C_{50} cyclic substituted alkylthiosemicarbazide, C_2-C_{20} straight chain alkenylthiosemicarbazide, C_2-C_{20} straight chain substituted alkenylthiosemicarbazide, C_3-C_{50} branched chain alkenylthiosemicarbazide, C_3-C_{50} branched chain substituted alkenylthiosemicarbazide, C_3-C_{50} cyclic alkenylthiosemicarbazide, C_3-C_{50} cyclic substituted alkenylthiosemicarbazide, C_2-C_{20} straight chain alkynylthiosemicarbazide, C_2-C_{20} straight chain substituted alkynylthiosemicarbazide, C_3-C_{50} branched chain alkynylthiosemicarbazide, C_3-C_{50} branched chain substituted alkynylthiosemicarbazide, C_3-C_{50} cyclic alkynylthiosemicarbazide, C_3-C_{50} cyclic substituted alkynylthiosemicarbazide, C_1-C_{20} straight chain alkylhydrazone, C_1-C_{20} straight chain substituted alkylhydrazone, C_2-C_{50} branched chain alkylhydrazone, C_2-C_{50} branched chain substituted alkylhydrazone, C_3-C_{50} cyclic hydrazoalkane, C_3-C_{50} cyclic substituted hydrazoalkane, C_2-C_{20} straight chain alkenylhydrazone, C_2-C_{20} straight chain substituted alkenylhydrazone, C_3-C_{50} branched chain alkenylhydrazone, C_3-C_{50} branched chain substituted alkenylhydrazone, C_3-C_{50} cyclic hydrazoalkene, C_3-C_{50} cyclic substituted hydrazoalkene, C_2-C_{20} straight chain alkynylhydrazone, C_2-C_{20} straight chain substituted alkynylhydrazone, C_3-C_{50} branched chain

- 17 -

alkynylhydrazone, C_3-C_{50} branched chain substituted
 alkynylhydrazone, C_3-C_{50} cyclic hydrazoalkyne, C_3-C_{50} cyclic
 substituted hydrazoalkyne, C_1-C_{20} straight chain alkylhydrazide,
 C_1-C_{20} straight chain substituted alkylhydrazide, C_3-C_{50} branched
 5 chain alkylhydrazide, C_3-C_{50} branched chain substituted
 alkylhydrazide, C_3-C_{50} cyclic alkylhydrazide, C_3-C_{50} cyclic
 substituted alkylhydrazide, C_2-C_{20} straight chain
 alkenylhydrazide, C_2-C_{20} straight chain substituted
 alkenylhydrazide, C_3-C_{50} branched chain alkenylhydrazide, C_3-C_{50}
 10 branched chain substituted alkenylhydrazide, C_3-C_{50} cyclic
 alkenylhydrazide, C_3-C_{50} cyclic substituted alkenylhydrazide, C_2-C_{20}
 straight chain alkynylhydrazide, C_2-C_{20} straight chain
 substituted alkynylhydrazide, C_3-C_{50} branched chain
 alkynylhydrazide, C_3-C_{50} branched chain substituted
 15 alkynylhydrazide, C_3-C_{50} cyclic alkynylhydrazide and C_3-C_{50}
 cyclic substituted alkynylhydrazide.

In preferred embodiments, polyamine species are linear
 or cyclic and are non-aromatic. In still more preferred
 embodiments, polyamine species are linear or cyclic, non-
 20 aromatic, and contain non-amide nitrogen atoms. By non-amide
 is meant a nitrogen which is not adjacent to a carbonyl group
 (i.e., $C=O$ or $C=S$).

In still other embodiments of the present invention
 compounds having the structure:




II

25 wherein R_4 is an oligonucleotide and M is a pendent group
 having an amine-containing species attached thereto are
 provided. The pendent group may be any group to which an
 amine-containing species may be attached. In preferred
 30 embodiments the pendent group is a $R_{10}S^-$ or $R_{10}NH^+$, wherein R_{10} is
 any of a broad range of reactive groups effective for

- 18 -

subsequent attachment of amine-containing species to the pendent group. Suitable for R_{10} are substituted and unsubstituted, straight chain or branched chained C_1 - C_{20} alkyl groups or substituted or unsubstituted C_7 - C_{14} aryl groups having the nucleophile in one position thereon and a further functional group in a further position thereon. The pendent group may thus, subsequently functionalized with a bifunctional linker group amendable for attachment of an amine-containing species to the pendent group. Alternatively the amine-containing species may be directly attached to a pendent group such as by alkylation.


Further in accordance with the present invention are provided compounds having the structure:



III

wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic amine-containing species containing non-amide nitrogen atoms, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle.

The present invention also provides novel amine containing compounds having the structure:



IV

wherein B is a purine or pyrimidine heterocycle, R_8 and R_9 are independently H, PO_2^- , a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_8 and R_9 is a purine containing oligonucleotide, and at least one of R_8 , R_9 and A is a species comprising the formula L_1 - L_2 -polyamine wherein L_1 is an amino linker and L_2 is a heterobifunctional linker; and wherein if R_8 is not a purine containing

- 19 -

oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- ; if R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH.


5 Thus R_8 and R_9 may be oligonucleotides and A may be a species comprising the formula L_1-L_2 -polyamine, or alternatively, R_8 may be an oligonucleotide and one or both of R_9 and A may be a species comprising the formula L_1-L_2 -polyamine; or R_9 may be an oligonucleotide and one or both of
10 R_8 and A may be a species comprising the formula L_1-L_2 -polyamine. Furthermore, when R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- . If R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide, and if A is
15 not a polyamine species then A is H or OH.

In preferred embodiments of the present invention commercially available amino linkers may be used. For example, the 3'-amino modifiers having the trade names C3 CPG and C7 CPG available through Glen Research may be employed. 5'-Amino
20 modifiers may also be used such as C3 and C7 5' branched modifiers available through Glen Research. Similarly, 2'-amino modifiers are also envisioned for use in some aspects of the present invention, see, e.g., United States Application Serial No. 782,374, filed 10/24/91. The amino linkers are designed to
25 functionalize a target oligonucleotide by the introduction of a primary amine at a designated site, be it 2', 3' or 5'. As will be apparent to one skilled in the art, any linker which meets this end is encompassed by the present invention.

Likewise, bifunctional linkers effective for purposes
30 of the present invention are available commercially. For example, bis-(maleimido)-methyl ether (BMME), disuccinimidyl suberate (DSS), 3-maleimidobenzoyl-N-hydroxy-succinimide (MBS), maleimidohexanoyl-N-hydroxyl-succinimide (MHS) and N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP) may be useful
35 in some embodiments of the present invention. Other useful bifunctional linkers will be apparent to one skilled in the art as for instance from Pierce, Rockford, IL.

- 20 -

Compounds of the present invention may be prepared by providing an oligonucleotide comprising one or more abasic sites. In the context of the present invention "abasic site" refers to a nucleotide unit in which the purine or pyrimidine group has been removed or replaced by a group such as a hydroxyl group. One or more abasic sites may be incorporated into one or more nucleotide bases of an oligonucleotide to form a synthon having the structure:



wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide, or amine-containing species, and at least one of R_1 and R_2 is a purine containing oligonucleotide, and X is H, $O-R_{11}$, $S-R_{11}$, F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkyaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide.

An enzymatic process may be used to produce such a synthon having abasic sites by reaction of a DNA glycosylase with an oligonucleotide starting material. Several glycosylase enzymes are available, see Friedberg, *DNA Repair* (W.H. Freeman

- 21 -

and Company, NY, 1985) p. 153. For example, uracil DNA glycosylase act on uracil bases within an oligonucleotide to create abasic sites. Of course, it should be recognized that enzymatic methods using DNA glycosylase may be less effective
5 for oligonucleotides more closely resembling RNA such as oligonucleotides having 2' modifications.


Enzymes, as employed in the present invention, may be derived from naturally occurring sources or may be prepared by recombinant techniques. Many useful enzymes are available
10 commercially.

Synthons alternatively can be prepared by incorporation of abasic sites into an oligonucleotide via abasic sugar precursors. For example, 5-O-(4,4'-dimethoxytrityl)-1,2-dideoxy-1-(o-nitrobenzyl)-D-ribofuranose-
15 3-O-(2-cyanoethyl-N,N'-diisopropyl) phosphoramidite may be prepared by modification of the procedures of Lyer, et al., *Nucleic Acids Research* 1990, 18, 2855 and Didier, et al., *Tetrahedron Letters* 1991, 32, 207. Phosphoramidites having a 2' substitutions and abasic sites may also be prepared. For
20 example, a synthon may have 2'-O-methyl or 2'-fluoro substitutions. Such phosphoramidite may be incorporated into an oligonucleotide by standard procedures. An o-nitrobenzyldeoxyfuranose containing oligonucleotide can be synthesized in accordance with these procedures. Post
25 synthesis photolysis utilizing a high intensity Hg lamp generates the corresponding abasic site-containing polymer. In addition, other methods of introducing abasic sites at the 3', 5' and internal positions of an oligonucleotide to form a synthon are known to those skilled in the art. Thereafter the
30 synthon may be reacted with an amine-containing species under reducing conditions. As illustrated in Figure 1, Step A, a compound may be prepared wherein B is uridine and an enzymatic process may be used to produce a synthon having abasic sites at one or more uridine sites by digestion of the compound with an
35 enzyme such as uracil-DNA glycosylase. Other glycosylases will be effective for different targets. As described above, a glycosylase may be determined by the combined sequence of R₁,

- 22 -

R_2 and B. Some useful glycosylases and their respective targets are described, for example, by Friedberg, *DNA Repair* (W.H. Freeman and Company, NY, 1985) p. 153. These enzymes are commercially available or may be prepared from known procedures in the art.

In other embodiments of the present invention, as exemplified in Figure 2, compounds having Formula II may be prepared by providing starting material having the structure:



wherein R_4 is an oligonucleotide, R_{12} is an oligonucleotide and B is urea or a heterocyclic base having a corresponding glycosylase and reacting the compound with an endonuclease to produce the compound 2 as described by Manoharan, et al., *J. Am. Chem. Soc.*, 1988, 110, 2690. Thereafter, the compound 2 is contacted with a pendent group such as $R_{10}S^+$, and reduced with the reducing agent $NaCNBH_4$ to stabilize the product 5. An amine-containing species may then be added such as by alkylation to provide the final product 7. An amine-containing species may alternatively be added directly to a bifunctional pendent group. Some endonucleases which will be useful in embodiments of the present invention are described, for example, in Doetsch et al., *Mutation Research* 1990, 236, 173, incorporated by reference herein in its entirety. The endonuclease chosen will depend upon the identity of B and the sequence of R_4 and/or R_{12} . Thus, if B is a pyrimidine heterocycle, and the sequence of R_{12} begins with a pyrimidine, then an endonuclease such as T4 or *M. luteus* UV endonuclease may be chosen. Following digestion by T4 or *M. luteus* UV endonuclease, B and R_{12} are removed, resulting in a 3' terminal α, β unsaturated aldehydic species. In some instances, it may be desirable to engineer the sequence of the species so as to provide a endonuclease digestion site at a desired location.

- 23 -

Thus, in one preferred embodiment of the present invention R_1 may be TGGGAGCCATAGCGAGGCUCG (SEQ ID NO: 1), B may be the pyrimidine thymine and R_{12} may be a thymidine dinucleotide. The net result of digestion of this species with
5 T4 UV endonuclease will be TGGGAGCCATAGCGAGGCN (SEQ ID NO: 2) wherein N represents the aldehydic species.

Treatment of the digested compound with pendent group comprising a linker bearing a nucleophile results in the addition of the pendent group at the 3' terminus of the
10 compound to join the linker to the digested compound. Suitable nucleophilic species include thiols and amines moieties as described above. In preferred embodiments of the present invention the pendent group is $R_{10}S^-$ or $R_{10}NH^-$. A polyamine species such as $NH_2(CH_2)_nNH_2$ wherein n is an integer from 1 to
15 about 10 could be used as the attacking nucleophile by suitably blocking one end thereof and utilizing the other end as the attacking nucleophilic species. R_{10} can be further selected to provide a linkage or bridge between the nucleophile and an amine-containing species. Suitable for R_{10} are substituted and
20 un-substituted, straight chain or branched chained C_1 - C_{20} alkyl groups or substituted or un-substituted C_7 - C_{14} aryl groups having the nucleophile in one position thereon and a further functional group in a further position thereon. After attachment of the pendent group via nucleophilic attack on
25 compound 2, for attachment of the amine-containing species the further functional group is then derivitized either via a bi-functional linking group, an alkylation type reaction or other derivation reaction known to those skilled in the art.

Upon addition of the pendent group to the digested
30 compound, the double bond remaining on the digested compound is reduced to stabilize the product. Reducing agents effective to stabilize the end product of such a reaction are well known in the art. Some suitable reducing agents include sodium cyanoborohydride, lithium cyanoborohydride and sodium
35 borohydride.

Thereafter an amine-containing species may be added via an alkylation reaction or directly to a pendent group which

- 24 -

is a bifunctional linker. The compound may further be derivatized by attaching one or more reactive groups to at least one of the nitrogen atoms of the amine-containing species. Reactive groups include, but are not limited to
5 reporter groups, alkylating agents, intercalating agents, RNA cleaving moieties, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins and cross-linking agents.

In accordance with other methods of the present invention compounds of Formula III may be prepared by reacting
10 an oligonucleotide having a 3' ribofuranosyl sugar with an oxidizing agent to produce an dialdehyde-terminated activated oligonucleotide. Suitable oxidants include periodate solution, lead tetraacetate, activated MnO_2 , thallium (III) salts, pyridinium chlorochromate and O_2 catalyzed by Co (III) salts.

15 Thereafter the dialdehyde-terminated activated oligonucleotide is reacted with an amine-containing species under reducing conditions. Reducing agents are known to those skilled in the art. Preferably, the activated oligonucleotide and species containing at least one nitrogen atom will be
20 reacted in the presence of a solution of sodium cyanoborohydride, lithium cyanoborohydride or sodium borohydride.

In preferred embodiments of the present invention compounds may be produced as illustrated by Figure 3, by
25 preparation of an oligonucleotide having a 3' ribofuranosyl end followed by attack of the 3' ribofuranosyl ring by an oxidant such as m-periodate solution in 0.1M NaOac buffer pH5, as described by Bayard, et al., *Biochemistry* 1986, 25, 3730 to produce a dialdehyde-terminated activated oligonucleotide
30 (Figure 3, Step A). The activated oligonucleotide and a species containing four nitrogen atoms, spermine, can be reacted in the presence of the reducing agent, sodium cyanoborohydride (Figure 3, Step B).

Compounds of the present invention preferably are
35 specifically hybridizable with a target region. By "specifically hybridizable" herein is meant capable of forming a stable duplex with a target DNA or RNA. It is believed that

- 25 -

oligonucleotides which form Watson-Crick base pairs, i.e., are complementary with target DNA or RNA and which specifically hybridize with target DNA or RNA, inhibit the flow of genetic information from DNA to protein. In some embodiments of the present invention the oligonucleotide portions of compounds of the present invention are at least 70% complementary to a target sequence. In preferred embodiments of the present invention the oligonucleotide portions of compounds of the present invention are at least 80% complementary to a target sequence. Full (100%) complementarity of the oligonucleotide portions of compounds of the present invention to a target sequence is most preferred. In preferred embodiments of the present invention, the oligonucleotide portions may be specifically hybridizable with DNA or RNA from papilloma virus, herpes viruses, human immunodeficiency virus, *Candida*, cytomegaloviruses, and influenza viruses. In addition, the oligonucleotide portions may also be specifically hybridizable with endogenous DNA or RNA of a cell. By oligonucleotide portions is meant R_1 and/or R_2 of Formula I, R_4 of Formula II, R_4 and/or R_6 of Formula III, or R_8 and/or R_9 of Formula IV. For therapeutics, an animal suspected of having a disease characterized by excessive or abnormal production of a protein is treated by administering a compound having the structure set forth in Formula I, Formula II, Formula III, or Formula IV in a pharmaceutically acceptable carrier. Most preferable, the compound is hybridizable with an RNA coding for the protein. Persons of ordinary skill in the art can easily determine optimum dosages, dosing methodologies and repetition rates. Such treatment is generally continued until either a cure is effected or a diminution in the diseased state is achieved. Long term treatment is likely for some diseases.

The compounds of the present invention will also be useful as a research reagent useful for the modulation of the production of a protein by an organism. Modulation may be accomplished by contacting the organism with compounds of the present invention having structures as set forth in Formula I,


- 26 -

Formula II, Formula III, or Formula IV. Preferably the compounds are hybridizable with RNA coding for the protein.

Diagnostic applications include the detection of the presence or absence of an RNA in a sample suspected of
5 containing RNA comprising contacting the sample with a compound having structures as set forth in Formula I, Formula II, Formula III or Formula IV wherein the compound is specifically hybridizable with the RNA and detecting the presence or absence of hybridization of the compound to the sample wherein
10 hybridization is indicative of the present of the RNA in the sample.

It is also envisioned by the present invention to provide compounds in which at least one of the nitrogen atoms of the polyamine are derivatized with one or more of the group
15 consisting of functionalities such as reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, crown amines, porphyrins, PNA (Peptide Nucleic Acids), PEG (polyethylene glycol) containing amines, amines and cross-linking agents. Therapeutic, diagnostic and
20 research reagent applications are equally, or even more effective when the polyamine species further comprises such groups. Such compounds allow greater numbers of functionalities to be delivered to a target. For example, reporter groups such as biotin, fluorescent molecules and
25 various fluorophores may be attached to compounds of the present invention to effect diagnostic ends, resulting in signal amplification as compared to conventional oligonucleotide-reporter group combinations. In a preferred embodiment of the present invention, biotin may be used to
30 functionalize compounds of the present invention by reacting a compound with D-biotin-N-hydroxysuccinimide ester. In a further preferred embodiment, the polyamine species may be further functionalized by reacting the compound containing the polyamine species with an activated ester having the structure
35 (Compound 13):

- 27 -



Where DNP stands for 2,4-dinitrophenyl protecting group to form a compound with repeating imidazole catalytic cleaver units useful as an antisense therapeutic agents. Heterobifunctional linkers also can be utilized for attachment of intercalators, RNA cleaving agents including imidazoles, cell receptor binding molecules, steroids, alkylating agents, crown amines, porphyrins and cross-linkers to the polyamine species.

The following examples are illustrative but are not meant to be limiting of the present invention.

10 EXAMPLE 1

Preparation of an Abasic Site Containing Oligonucleotide via Enzymatic Reaction

A. Synthesis of an Oligonucleotide containing a Single Uridine Site

15 An oligonucleotide having the sequence CGC AGU CAG CC (SEQ ID NO:3) wherein U represents a 2'-deoxyuridine nucleotide, was prepared by standard solid phase synthesis. The deoxyuridine nucleotide in the middle of the sequence was added during synthesis utilizing deoxyuridine phosphoramidite (Glen Research, Sterling, VA). The oligonucleotide was prepared
20 utilizing standard synthesis cycles. It was deprotected by normal deprotection at 55°C utilizing ammonium hydroxide, 30%, for 16 hours. The solvent was evaporated and the residue was purified by HPLC and detritylated. Final purification was
25 effected on Sephadex G-25.

B. Preparation of Enzyme Stock Solution

Uracil-DNA glycosylase was isolated from *E. Coli* M5219 cells transformed with the expression plasmid pBD396 containing

- 28 -

the *ung* gene. The enzyme was purified by electrophoretic homogeneity as described by Lindahl, et al., *J. Biol. Chem.* 1977, 252, 3286 and stored in 30mM HEPES-NaOH, pH 7.4, containing 5% glycerol, 2mM DTT and 1 mM EDTA.

5 C. **Preparation of Oligonucleotide Containing Single Abasic Site**

 An abasic oligonucleotide of the sequence CGC AGN CAG CC (SEQ ID NO:4) wherein N represents an abasic site, was prepared by treating 237 O.D. units of an
10 oligonucleotide having SEQ ID NO:1 of Example 1A in 0.5ml water with 200 μ l of the stock solution of Example 1B (200 μ g of uracil DNA-glycosylase) and incubating at room temperature overnight. HPLC analysis showed quantitative removal of uracil as indicated by a 1:10 ratio between uracil and the abasic
15 dodecamer oligonucleotide. The uracil retention time was 2.43 minutes and the abasic oligonucleotide was 21.68 minutes. The solution was lyophilized and stored in the freezer until further use.

20 D. **Preparation of Oligonucleotide Containing Multiple Uridine Sites**

 In the manner of Example 1A the following oligonucleotide was prepared GAC AGA GGU AGG AGA AGU GA (SEQ ID NO: 5) wherein U represents a 2'-deoxyuridine nucleotide. The oligonucleotide is treated in accordance with the procedure of
25 Example 1C resulting in an oligonucleotide of the sequence GAC AGA GGN AGG AGA AGN GA (SEQ ID NO: 6) wherein N represents an abasic site within the oligonucleotide.

EXAMPLE 2

30 **Preparation of an Abasic Site Containing Oligonucleotide via an Abasic Sugar Precursor**

 A. **Preparation of 5-O-4,4'-Dimethoxytrityl-1,2-Dideoxy-1-(o-nitrobenzyl)-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.**

 5-O-4,4'-dimethoxytrityl-1,2-dideoxy-D-ribofuranose-3-
35 O-(2-cyanoethyl-N,N'-diisopropyl) phosphoramidite is prepared in accordance with modification of the procedures of Lyer, et

- 29 -

al., *Nucleic Acids Research* 1990, 18, 2855 and Didier, et al., *Tetrahedron Letters* 1991, 32, 207 incorporated by reference herein in their entireties.

5 B. **Preparation of Oligonucleotide Containing Abasic Site**

Oligonucleotide having the sequence CGC AGN CAG CC wherein N represents an abasic site (SEQ ID NO:4) from Example 1C can be prepared in accordance with modifications of the procedures of Lyer, et al., *Nucleic Acids Research* 1990, 18, 10 2855 and Didier, et al., *Tetrahedron Letters* 1991, 32, 207. In accordance with these procedures, an o-nitrobenzyl deoxyfuranose containing oligonucleotide is synthesized using the oligonucleotide synthetic methods of Lyer, et al., and Didier, et al., Photolysis utilizing a high intensity Hg lamp 15 (300nm) generates the corresponding abasic site containing oligonucleotide. Such abasic oligonucleotides are also described in Horn, et al., *Nucleosides and Nucleotides* 10:299 (1991).

EXAMPLE 3

20 **Preparation of Modified Abasic Sugar Precursors**

A. **Preparation of 5-O-(4,4'-Dimethoxytrityl)-2-O-Methyl-1,2-dideoxy-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.**

1-O-methyl-D-ribofuranose is 3,5 protected with TIPS- 25 Cl₂. It is then 2-position methylated with either diazomethane or methyl iodide/silver oxide (CH₃I/Ag₂O). The composition is then treated with an acetic anhydride/acetic acid/sulfuric acid mixture to give a 1-O-acetyl, 2-O-methyl 3,5 protected sugar. The 1-O-acetyl, 2-O-methyl 3,5 protected sugar is deprotected 30 with tetrabutyl ammonium fluoride, 5-position dimethoxytritylated, and 3-position phosphitylated. Thereafter, this phosphoramidite may be incorporated into an oligonucleotide by standard phosphoramidite procedures and ammonia deprotected to form a 2'-O-methyl, 1' abasic site 35 containing oligonucleotide.

- 30 -

B. Preparation of 5-O-(4,4'-Dimethoxytrityl)-2-O-Methyl-1,2-Dideoxy-1-(o-nitrobenzyl)-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.

5 1-O-acetyl 2,3,5-tri-O-benzoyl-D-ribofuranose is condensed with o-nitrobenzyl alcohol under Vorbruggen conditions. The resultant 1-O-(ortho-nitrobenzyl)-2,3,5-tri-O-benzoyl (α,β)-D-ribofuranose is deprotected with ammonia and subsequently treated with TIPS-Cl₂. The resultant 3,5-silyl
10 protected 1-O-(ortho-nitrobenzyl) D-ribofuranose is reacted with diazomethane or CH₃I/Ag₂O to give the required 2-O-methyl compound. Subsequent 3,5-deprotection, 5-dimethoxy tritylation and 3-phosphitylation gives the named phosphoramidite. The phosphoramidite can be incorporated into an oligonucleotide via
15 standard phosphoramidite procedures.

C. Preparation of 5-O-(4,4'-Dimethoxytrityl)-2-Fluoro-1,2-Dideoxy-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.

1-O-(ortho-nitrobenzyl)-2,3,5-tri-O-benzoyl-D-
20 ribofuranose is deprotected at 2,3,5 positions using ammonia. Tritylation with excess trityl chloride/pyridine/4-dimethylaminopyridine gives 3-5-ditrityl-1-O-nitrobenzyl-D-ribofuranose. Oxidation at 2 position with CrO₃ followed by NaBH₄ reduction inverts the configuration at 2 position yielding an
25 arabino sugar. The arabino sugar is converted to its triflate at 2 position and the triflate is displaced with fluoride ion to yield the 2-fluoride modified sugar which can be 5 position protected and phosphitylated to incorporate the sugar into an oligonucleotide via standard oligonucleotide synthesis.

- 31 -

EXAMPLE 4

Oligonucleotides conjugated in the following example are set forth in Table 2.

TABLE II

5	OLIGOMER (SEQ ID NO.)	TARGET	SEQUENCE	LINKER (L)	OTHER MODIFI- CATIONS
	A (SEQ ID NO:7)	ICAM	TGG GAG CCA TAG CGA GGC-L	3-carbon amino	P=S
10	B (SEQ ID NO:7)	ICAM	TGG GAG CCA TAG CGA GGC-L	3-carbon amino	P=O
	C (SEQ ID NO:8)	BPV	CTG TCT CCA* TCC TCT TCA CT	2'-amino- pentoxy	P=O
	D (SEQ ID NO:9)	BPV	CTG TCT CCA TCC TCT TCA CT-L	3-carbon amino	P=O
15	E (SEQ ID NO:9)	BPV	CTG TCT CCA TCC TCT TCA CT-L	6-carbon amino	P=O
	F (SEQ ID NO:10)	CMV	GGC GUC UCC AGG CGA UCU GAC*		2'-OMe
20	G (SEQ ID NO:11)	ICAM	TCT GAG TAG CAG AGG AGC TC*		2'-OMe
	H (SEQ ID NO:12)		GGA UGG CGU CUC CAG GCG AUC*		2'-OMe
	I (SEQ ID NO:13)		GGA UGG CGU CUC CAG GCG AUC-L	3-carbon amino	2'-OMe
25	J (SEQ ID NO:13)		GGA UGG CGU CUC CAG GCG AUC-L	6-carbon amino	2'-OMe
	K (SEQ ID NO:7)		F-TGG GAG CCA TAG CGA GGC-L	3-carbon amino	2'-OMe

A = 2'-O-aminopentoxy-2'-deoxyadenosine

30 C = 2'-aminopropoxy cytosine

F = Fluorescein

- 32 -

A. 3' Terminus Polyamine End Labeled Oligonucleotide.

1. 3'-Terminus Polyamine Oligonucleotide I

Polyamines were attached to the 3'-terminus end of a
5 phosphodiester oligonucleotide having the sequence D-polyamine
[(SEQ ID NO: 9)-polyamine], wherein the polyamine is one of the
following:

TABLE III

	- 1,6 Diaminohexane	Oligomer D(i)
10	- Diethylenetriamine	Oligomer D(ii)
	- Triethylenetetramine	Oligomer D(iii)
	- Spermine	Oligomer D(iv)
	- Pentaethylenhexamine	Oligomer D(v)

a. Preparation of the Intermediate Linker

15 The oligonucleotide sequence having a 3'-terminus
amino group was synthesized using 3'-amino modifier (with a
three carbon linker) controlled pore glass (CPG) from Glen
Research as the solid support. The synthesis was conducted
with an Applied Biosystems 380B or 994 in the "Trityl-Off"
20 mode. The resultant oligonucleotide was cleaved from the solid
support and deprotected with concentrated NH_4OH for 16 hrs at
55°C. Purification on a Sephadex G-25 column yielded a 3'-
amino modified oligonucleotide of the specified sequence.

b. Preparation of Polyamine Functionalized Oligonucleotide

25 The crude 3'-aminolinker-oligonucleotide (SEQ ID NO:9)
(15 O.D. units, approximately 85 nmols) was dissolved in
freshly prepared NaHCO_3 buffer (150 ul, 0.2 M, pH 8.1) and
treated with a solution of disuccinimidyl suberate (DSS)
30 (approximately 5 mgs) dissolved in 150 ul of methyl sulfoxide
(DMSO). The reaction mixture was left to react for 20 minutes
at room temperature. The mixture was then passed over a
Sephadex G-25 column (0.7 x 45 cm) to separate the activated
oligonucleotide-DSS from the excess DSS. The oligonucleotide-
35 DSS was then frozen immediately and lyophilized to dryness. A

- 33 -

solution of polyamine in 0.33 M NaOAc (approximately 6 mg polyamine in 300 ul 0.33 M NaOAc, pH 5.2, final solution pH 6-8.0) was added to the dried oligonucleotide-DSS, and this mixture was allowed to react overnight at room temperature.

- 5 The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using
- 10 a Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 4.

TABLE IV

	Oligomer	Retention Time
	unreacted D	26.44 mins
15	Oligomer D(i)	27.48 mins
	Oligomer D(ii)	27.23 mins
	Oligomer D(iii)	27.27 mins
	Oligomer D(iv)	27.54 mins
	Oligomer D(v)	27.36 mins

- 20 In a second test run under the same conditions the HPLC gradient was 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 15% solvent B in 60 minutes. HPLC retention times were as set forth in Table 5.

TABLE V

25	Oligomer	Retention Time
	untreated D	60.74 mins
	Oligomer D(ii)	62.37 mins
	Oligomer D(v)	65.24 mins

- Gel analysis showed progressively slower migration
- 30 times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Gel: 313-107)

- 34 -

c. Nuclease stability of 3' polyamine
conjugates in Fetal Calf Serum

Polyamine conjugates of the invention are assessed for their resistance to serum nucleases by incubation of the oligonucleotides in media containing various concentrations of fetal calf serum. Labeled oligonucleotides are incubated for various times, treated with protease K and then analyzed by gel electrophoresis on 20% polyacrylamide-urea denaturing gels and subsequent autoradiography or phosphor-imaging. Autoradiograms are quantitated by laser densitometry. Based upon the location of the modifications and the known length of the oligonucleotide it is possible to determine the effect of the particular modification on nuclease degradation. For the cytoplasmic nucleases, a HL60 cell line is used. A post-mitochondrial supernatant is prepared by differential centrifugation and the labeled oligonucleotides are incubated in this supernatant for various times. Following the incubation, oligonucleotides are assessed for degradation as outlined above for serum nucleolytic degradation. Autoradiography results are quantitated for comparison of the unmodified and the modified oligonucleotides.

The $t_{1/2}$ are set forth below:

TABLE VI

	Oligonucleotide	$t_{1/2}$ (hours)
25	wild type oligomer D	0.5 (no aminolinker)
	unreacted oligomer D	22 (with aminolinker)
	oligomer D(ii)	48
	oligomer D(v)	>50

2. 3' - Terminus Polyamine Conjugate II

30 Polyamines were attached to the 3'-terminus end of a phosphodiester oligonucleotide having the sequence E-polyamine

- 35 -

[(SEQ ID NO: 9)-polyamine] wherein the polyamine is one of the following:

TABLE VII

- 5 - Diethylenetriamine Oligomer E(i)
 - Pentaethylenhexamine Oligomer E(ii)

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a substituting a 3' amino modifier with a six carbon linker (Clonetech, Palo Alto, CA) for the 3'-amino
10 modifier (with a three carbon linker).

3. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide was prepared in accordance with Example 4-A-1-b. The resulting
15 polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 25% solvent B in the next 50 minutes using a Water's Delta-Pak C-18
20 reverse phase column. HPLC retention times were as set forth in Table 8.

TABLE VIII

	Oligomer	Retention Time
	untreated E	41.38 mins
25	Oligomer E(i)	43.29 mins
	Oligomer E(ii)	43.43 mins

Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone.
30 (Gel: 353-35).

4. 3' - Terminus Polyamine Conjugate III

Polyamines were attached to the 3'-terminus end of a phosphorothioate oligonucleotide having the sequence A-

- 36 -

polyamine [(SEQ ID NO:7)-polyamine] where the polyamine is one of the following:

TABLE IX

	- 1,6 Diaminohexane	Oligomer A(i)
5	- Diethylenetriamine	Oligomer A(ii)
	- Triethylenetetramine	Oligomer A(iii)
	- Spermine	Oligomer A(iv)
	- Pentaethylenehexamine	Oligomer A(v)

a. Preparation of the Intermediate Linker

10 The intermediate linker was prepared as described in Example 4-A-1-a utilizing the Beaucage reagent (3H-1,2-benzodithioate-3-one 1,1-dioxide, Radhakrishnan, et al., *J. Am. Chem. Soc.* 1990, 112, 1253) to form the phosphorothioate internucleotide backbone. The 3'-aminolinker was introduced as
15 described in example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

Oligonucleotides were functionalized as described in Example 4-A-1-b. The resulting polyamine-oligonucleotide
20 conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using a Water's Delta-Pak C-18 reverse phase column. HPLC
25 retention times were as set forth in Table X.

TABLE X

	Oligomer	Retention Time
	unreacted A	30.77 mins
	Oligomer A(iii)	31.31 mins
30	Oligomer A(v)	31.02 mins

In a second test run under the same conditions, the HPLC gradient was 0-10 mins, 95% solvent A, 5% solvent B;

- 37 -

linear increase to 15% solvent B in 60 minutes. Retention times were as set forth in Table XI.

TABLE XI

	Oligomer	Retention Time
5	untreated A	68.62 mins
	Oligomer A(i)	68.70 mins
	Oligomer A(ii)	68.69 mins

In a second test run under the same conditions, HPLC retention times were as set forth in Table XII.

10

TABLE XII

	Oligomer	Retention Time
	untreated A	30.34 mins
	Oligomer A(iv)	30.57 mins
	Oligomer A(v)	30.72 mins

15

Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Test run 1 Gel, 313-82; Test run 2 Gel, 285-138; Test run 3 Gel, 353-57)

20

c. Preparation of Biotin Functionalized Oligonucleotide Polyamine Conjugate

To further characterize the oligonucleotide polyamine conjugate, biotin was attached to the free amines made available by the polyamines attached in Example 4-A-4-b. About
25 10 O.D. units (A_{260}) of Oligomers A(i) and A(ii) (approximately 58 nmoles) were dried in a microfuge tube. The oligonucleotide polyamine conjugate was rehydrated in 400 ul of 0.2 M NaHCO_3 (pH 8.1) buffer and D-biotin-N-hydroxysuccinimide ester (approximately 5.0 mgs biotin for the 1,6 Diaminohexane
30 conjugate, 8.0 mgs for the Diethylenetriamine) (Sigma) was added followed by 200 ul of DMF. The solution was left to react overnight at room temperature. The solution was then passed over a NAP-25 column and analyzed by reverse phase HPLC. Solvent A was 50 mM TEAA and solvent B was CH_3CN . The HPLC
35 gradient was 0-10 mins, 95% A, 5% B; linear increase to 40% B in the next 50 minutes using a Water's Delta-Pak C-18,

- 38 -

reverse phase column. The HPLC retention times were as set forth in Table XIII.

TABLE XIII

	Oligomer	Retention Time
5	untreated A	30.77 mins
	Oligomer A(i)	31.31 mins
	Oligomer A(i)-Biotin	35.56 mins
	Oligomer A(ii)	31.02 mins
	Oligomer A(ii)-Biotin	36.23 mins

10

5. 3' - Terminus Polyamine Conjugate IV

Polyamines were attached to the 3'-terminus end of the phosphodiester oligonucleotide having the sequence B-polyamine [(SEQ ID NO: 7)-polyamine] wherein the polyamine is one of the following:

15

TABLE XIV

- Diethylenetriamine	Oligomer B(i)
- Triethylenetetramine	Oligomer B(ii)
- Spermine	Oligomer B(iii)
- Pentaethylenhexamine	Oligomer B(iv)

20

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

25

The oligonucleotide was functionalized with polyamines as described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in 30 the next 50 minutes using a Water's Delta-Pak C-18 reverse

- 39 -

phase column. HPLC retention times were as set forth in Table XV.

TABLE XV

	Oligomer	Retention Time
5	untreated B	25.71 mins
	Oligomer B(i)	26.11 mins
	Oligomer B(ii)	25.26 mins
	Oligomer B(iii)	25.10 mins
	Oligomer B(iv)	25.12 mins
10	Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Gel: 313-112)	

B. 2' Internal Polyamine Labeled Oligonucleotide

- 15 **1. 2'-Internal Polyamine Oligonucleotide I**
- Polyamines were attached to the 2'-internal linker site of a phosphodiester oligonucleotide having the sequence C-polyamine [(SEQ ID NO: 8)-polyamine] wherein the polyamine is one of the following:

20 TABLE XVI

- Diethylenetriamine	Oligomer C(i)
- Triethylenetetramine	Oligomer C(ii)
- Pentaethylenhexamine	Oligomer C(iii)

a. Preparation of the Intermediate Linker

- 25 The intermediate linker was prepared as described in Example 4-A-1-a incorporating a modified adenosine phosphoramidite (with a 2'-aminolinker) at position #9. This oligonucleotide and the 2'-amino linker have been described in Manoharan, et al., *Tetrahedron Letters* 1991, 32, 7171.

30 **b. Preparation of Polyamine Functionalized Oligonucleotide**

- The oligonucleotide was functionalized as described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent

- 40 -

B; linear increase to 40% solvent B in the next 50 minutes using a Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XVII.

TABLE XVII

5	Oligomer	Retention Time
	untreated C	26.20 mins
	Oligomer C(i)	27.52 mins
	Oligomer C(ii)	27.50 mins
	Oligomer C(iii)	27.59 mins
10	Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Gel: 313-97)	
15	C. 3' Terminus Polyamine End Labeled Oligonucleotide, Using a 2'-aminolinker	
	1. 3' Terminus Polyamine Labeled Oligonucleotide I, Using a 2'-aminolinker	

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide via a 2'-aminolinker having the sequence F-polyamine [(SEQ ID NO:10)-polyamine] wherein the polyamine is pentaethylenehexamine (oligomer F(i)).

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a, except that a modified cytosine CPG (with a 2'-propylaminolinker) was introduced at the 3' end. The 2'-modification can be prepared by modification of the procedure previously described in Application Serial No. 918,362 filed July 23, 1992. The CPG containing 2'- ω -phthalimido-propoxy-cytidine was synthesized according to the standard protocols reported in the literature. See, for example, B. S. Sproat and A.I. Lamond, in "Oligonucleotides and Analogues" edited by F. Eckstein, IRL Press at Oxford University Press (1991) p71-72.

b. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide was prepared in accordance with Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by

- 41 -

reverse phase HPLC and a 20% denaturing gel. Solvent A was 50mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XVIII.

TABLE XVIII

	Oligomer	Retention Time
	unreacted F	28.53 mins
10	oligomer F(i)	29.47 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

15 2. 3' Terminus Polyamine Labeled
Oligonucleotide II, Using a 2'-aminolinker

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide via a 2'- aminolinker having the sequence G-polyamine [(SEQ ID NO:11)-polyamine] wherein the polyamine is pentaethylenehexamine (oligomer G(i)).

20 a. Preparation of the Intermediate Linker

The intermediate linker was prepared in accordance with the method described in Example 4-A-1-a.

 b. Preparation of Polyamine
 Functionalized Oligonucleotide

25 The polyamine functionalized oligonucleotide was prepared in accordance with the procedures described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50mM TEAA, solvent B was CH₃CN. The HPLC
30 gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XIX.

- 42 -

TABLE XIX

Oligomer	Retention Time
unreacted G	28.43 mins
oligomer G(i)	29.06 mins

5 Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

3. 3' Terminus Polyamine Labeled
Oligonucleotide III Using a 2'-aminolinker

10 Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide via a 2'-aminolinker having the sequence H-polyamine [(SEQ ID NO:12)-polyamine] wherein the polyamine is pentaethylenehexamine.

a. Preparation of the Intermediate Linker

15 The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

b. Preparation of Polyamine
Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is
20 prepared in accordance with methods described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B;
25 linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XX.

TABLE XX

Oligomer	Retention Time
unreacted H	28.49 mins
oligomer H(i)	30.36 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

35 EXAMPLE 5

- 43 -

Polyamine Labeled 2'-OMe Oligonucleotides and Other RNA Mimics**A. Polyamine Labeled 2'-OMe Oligonucleotide I**

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide (via a 3 carbon linker) having the sequence I-polyamine [(SEQ ID NO:13)-polyamine] wherein the polyamine is pentaethylenehexamine (oligomer I(i)).

1. Preparation of the Intermediate Linker

The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

2. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is prepared in accordance with methods described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XXI.

TABLE XXI

Oligomer	Retention Time
unreacted I	28.93 mins
oligomer I(i)	29.59 mins
Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 353-156)	

B. Polyamine Labeled 2'-OMe Oligonucleotide II

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide (via a 6 carbon linker) having the sequence J-polyamine [(SEQ ID NO:13)-polyamine] wherein the polyamine is pentaethylenehexamine (oligomer J(i)).

1. Preparation of the Intermediate Linker

The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

- 44 -

2. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is prepared in accordance with methods described in Example 4-A-1-
 5 b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using
 10 Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XXII.

TABLE XXII

	Oligomer	Retention Time
	unreacted J	28.76 mins
15	oligomer J(i)	29.39 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

C. Polyamine Labeled 2'-OMe Oligonucleotide III

20 Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide (via a 3 carbon linker) having another reporter group (such as biotin, fluorescein) at the other end in the sequence K-polyamine [(SEQ ID NO:7)-polyamine]. Fluorescein at 5' end was added using the required
 25 amidite commercially available from Clontech. The polyamine is one of the following

- pentaethylenhexamine oligomer K(i)
- spermine oligomer K(ii)

1. Preparation of the Intermediate Linker

30 The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

2. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is
 35 prepared in accordance with methods described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel.

- 45 -

Solvent A was 50mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XXIII.

TABLE XXIII

	Oligomer	Retention Time
	unreacted K	31.35 mins
	oligomer K(i)	31.96 mins
10	oligomer K(ii)	32.15 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 353-149)

EXAMPLE 615 **5' Terminus Polyamine End Labeled Oligonucleotide****A. 5'-Terminus Polyamine Oligonucleotide I**

Polyamines were attached to the 5'-terminus end of a phosphodiester oligonucleotide having the following sequences:

- 5'-aminolinker-TCAG (oligomer L)
- 20 - 5'-aminolinker-CGCACGC (oligomer M)

to provide the polyamine oligonucleotides:

- 5'-polyamine-TCAG (oligomer L(i))
- 5'-polyamine-CGCACGC (oligomer M(i)) wherein the polyamine is pentaethylenehexamine.

25 **1. Preparation of the Intermediate Linker**

The oligonucleotide sequence having a 5'-terminus amino group was synthesized using Aminolink-II (with a six carbon linker) phosphoramidite from Applied Biosystems in the last round of synthesis. The synthesis was conducted with an Applied Biosystems 380B or 994 in the "Trityl-On" mode. The resultant oligonucleotide was cleaved from the solid support and deprotected with concentrated NH₄OH for 16 hrs at 55° C. Purification on a Sephadex G-25 column yielded a 5'-amino modified oligonucleotide of the specified sequence.

- 46 -

2. Preparation of Polyamine Functionalized Oligonucleotide L(i)

The crude 5'-aminolinker-oligonucleotide (150 O.D. units, approximately 3.75 mmols) was dissolved in freshly prepared NaHCO₃ buffer (900 ul, 0.2 M, pH 8.1) and treated with a solution of disuccinimidyl suberate (DSS) (approximately 30 mgs) dissolved in 750 ul of methyl sulfoxide (DMSO). The reaction mixture was left to react for 20 minutes at room temperature. The mixture was divided into three portions and then passed over a Sephadex G-25 column (0.7 x 45 cmx3 columns) to separate the activated oligonucleotide-DSS from the excess DSS. The oligonucleotide-DSS was then frozen immediately and lyophilized to dryness. A solution of polyamine in 0.33 M NaOAc (approximately 60 mL polyamine in 1950ul 0.33 M NaOAc, pH 5.2, final solution pH 6-8.0) was added to the dried oligonucleotide-DSS, and this mixture was allowed to react overnight at room temperature.

3. Preparation of Polyamine Functionalized Oligonucleotide M(i)

The crude 5'-aminolinker-oligonucleotide (oligomer M) (150 O.D. units, approximately 2.50) was reacted as described in Example 6(b).

4. Characterization of 5' Polyamin Functionalized Oligonucleotides

The resulting polyamine-oligonucleotide conjugates were characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table XXIV.

TABLE XXIV

	Oligomer	Retention Time
	unreacted L	22.78 mins
35	oligomer L(i)	28.27 mins
	unreacted M	24.50 mins
	oligomer M(i)	26.72 mins

- 47 -

EXAMPLE 7**Preparation of a Reactive Site containing Oligonucleotide**

An oligonucleotide having the sequence TGGGAGCCATAGCGAGGUCT (SEQ ID NO: 14) is treated with uracil DNA
5 glycosylase followed by T4 endonuclease. The product is then treated with 1-phthalimidobutyl-4-thiol. Nucleophilic attack by the thiol with the protected aminobutyl moiety results in addition to what was the 3' position of the opened nucleotide. Treatment of this composition with hydrazine will deblock the
10 phthalimide yielding an amino species which is then treated with bifunctional linker followed by treatment with an appropriate polyamine species as per Example 4-A-1-b.

EXAMPLE 8**Preparation of Polyamine Conjugated Oligonucleotide**

15 An oligonucleotide is prepared as described in Example 7 treating the product with $\text{NH}_2\text{-CH}_2\text{-CH}_2\text{-SH}$. The thiol group will attack the double bond of the opened nucleotide. The resulting amine may then be further derivatized with a reactive group.

20 EXAMPLE 9**Thermodynamic Parameters of Oligoamine-Oligonucleotide Conjugates with DNA and RNA Targets**

The ability of the functionalized oligonucleotides of the invention to hybridize to their complementary RNA or DNA
25 sequences is determined by thermal melting analysis. The RNA complement is synthesized from T7 RNA polymerase and a template-promoter of DNA synthesized with an Applied Biosystems, Inc. 380B nucleic acid synthesizer. The RNA species is purified by ion exchange using FPLC (LKB Pharmacia,
30 Inc.) or by denaturing urea-PAGE. Natural antisense oligonucleotides or those containing functionalization at specific locations are added to either the RNA or DNA complement at stoichiometric concentrations to form hybrid duplexes. The absorbance (260 nm) hyperchromicity dependence
35 on temperature upon duplex to random coil transition is

- 48 -

monitored using a Gilford Response II spectrophotometer. These measurements are performed in a buffer of 10 mM Na-phosphate, pH 7.4, 0.1 mM EDTA, and NaCl to yield an ionic strength of either 0.1 M or 1.0 M. Data are analyzed by a graphic
5 representation of $1/T_m$ vs $\ln[Ct]$, where $[Ct]$ is the total oligonucleotide concentration. From this analysis the thermodynamic parameters are determined. Based upon the information gained concerning the stability of the duplex or hetero-duplex formed, the placement of the polyamines into
10 oligonucleotides is assessed for its effects on helix stability. Modifications that drastically alter the stability of the hybrid exhibit reductions or enhancements in the free energy (ΔG) and decisions concerning their usefulness in antisense oligonucleotides are made.

TABLE XXV

Oligomer	DNA TARGET			RNA TARGET		
	T _m (°C)	ΔT _m (°C)	ΔΔG° _{37°C}	T _m (°C)	ΔT _m (°C)	ΔΔG° _{37°C}
wild type oligomer D	60.6	-	-	64.9	-	-
5 oligomer D	60.3	-0.3	+0.3	64.6	-0.3	0.0
oligomer D + 5'- 6-carbon amino linker)	60.8	+0.2	0.0	65.1	+0.2	0.0
oligomer E	60.8	+0.2	-0.8	65.8	+0.9	-1.0
oligomer E(i)	61.2	+0.6	-1.4	66.3	+1.4	-1.9
10 oligomer E + spermine	61.5	+0.9	-1.7	67.1	+2.2	-2.1
oligomer E(ii)	61.2	+0.6	-1.3	67.5	+2.6	-2.6

- 50 -

EXAMPLE 10**Conjugation of Polyamines to Abasic Site-Containing Oligonucleotides**

To 15.2 ODS of an abasic oligonucleotide (SEQ ID NO: 5 4) in 100 μ l water was added 25 μ l 1M NaOAc (pH 5.0) solution. The final concentration of the acetate buffer was 0.2 M. 5.3 mg of triethylenetetramine was dissolved in 500 μ l of 1M NaOAc (pH 5.0) solution. 50 μ l of the resulting solution was added to the oligonucleotide solution followed by 50 μ l of NaCNBH₃ 10 (57 MM solution). The pH of the resulting solution was below 8.0. The solution was vortexed and left to stand overnight. HPLC and Gel analysis indicated conjugation of the triethylenetetramine to the oligonucleotide. The conjugated oligonucleotide was purified by G-25 and HPLC. HPLC retention 15 times are set forth in Table XXVI.

TABLE XXVI

	Oligomer	Retention time (mins)
	parent oligonucleotide (SEQ ID NO:3)	26.66
20	abasic oligonucleotide (SEQ ID NO:4)	26.16
	(SEQ ID NO:4)-triethylenetetramine conjugate	26.04

EXAMPLE 11**25 Oligonucleotide Synthesis**

Oligonucleotides of the following sequences were synthesized with an Applied Biosystems 380B or 994 in the "Trityl On" mode. The resultant oligonucleotides were cleaved from the solid support and deprotected with concentrated NH₄OH 30 for 16 hour at 55°C. HPLC purification with a Water's Delta-Pak C-18, reverse phase column followed with the given gradient: Solvent A: 50 mM TEAA, pH=7.4; Solvent B: CH₃CN; 0-10 mins., 95% A, 5% B; linear increase to 60% B in the next fifty mins. The full-length, DMT-on oligonucleotide was 35 separated from the impurities. Treatment with 80% acetic acid

- 51 -

removed the DMT. A final run over a Sephadex G-25 column yielded pure oligonucleotides of the specified sequences.

CGC AGU CAG CC (SEQ ID NO:3)

GAU CT (SEQ ID NO:15)

5 EXAMPLE 12

Abasic Site Generation

To generate an abasic site at the uracil position in the sequences prepared in Example 11, uracil DNA glycosylase was added to the oligonucleotides (approximate
10 ratio 100 O.D. oligonucleotide to 100 "units" enzyme). This was left to react overnight at room temperature. HPLC analysis (HPLC gradient was as follows: Solvent A: 50 mM TEAA, pH=7.4; Solvent B: CH₃CN; 0-10 mins., 95% A, 5% B; linear increase to 15% B in the next fifty mins. HPLC column: Water's
15 Delta-Pak C-18, reverse phase) shows a small excess uracil peak at 2.58 minutes and the oligonucleotides with the abasic site at 33.38 minutes.

CGC AGN CAG CC (SEQ ID NO:4)

GAN CT (SEQ ID NO:16)

20 (N = abasic site)

EXAMPLE 13

Conjugation To Oligonucleotides Containing Abasic Sites

A. Oligonucleotide Having SEQ ID NO:4.

The oligonucleotide having sequence CGC AGN CAG CC
25 (SEQ ID NO:4) was divided into 25 O.D. unit samples for conjugation. A 50 µl portion of 1M NaOAc was added to each of these samples (25 O.D./100 µl of HPLC grade water) to assure a low pH. The following solutions were made:

TABLE XXVII

Ligand	mg/ μ l (pH=5.0)	1M NaOAc	DMF
6-((biotinoyl)amino) 5 caproic acid hydrazide	5 mg	300 μ l	100 μ l
fluorescein-5- thiosemicarbazide	5 mg	200 μ l	100 μ l
Lys-Tyr-Lys(tripeptide)	5 mg	200 μ l	
Lys-Trp-Lys(tripeptide)	5 mg	200 μ l	
10 triethylenetetramine (TEA)	5 mg	200 μ l	
pentaethylenhexamine (PEHA)	5 mg	200 μ l	
5-amino-O-phenanthroline	5 mg	200 μ l	100 μ l
15 1-pyrene-butryl- hydrazide	5 mg	200 μ l	100 μ l
PEG-hydrazide (methoxy polyethylene glycol- carboxymethyl hydrazide)	5 mg	200 μ l	

20 A 100 μ l portion of each of the solutions given in Table XXVII was added to the oligonucleotide solutions; 5 mgs of PEG-hydrazide were added directly. After about period of 15 minutes, 100 μ l of a NaCNBH₃ solution (0.20 M NaBH₃CN in 0.25 M NaOAc) was added to each of the reactions. The 25 reaction mixtures were then put on a vortex-shaker and left overnight at room temperature. The conjugates were then analyzed by HPLC and 20% PAGE gel, indicating formation of the conjugate. The results are shown in Table XXVIII.

- 53 -

TABLE XXVIII

Oligo./Conjugate	HPLC Program	Retention Time (min.)	Yield (O.D.)
SEQ ID NO:3	biohr	33.09	
5 SEQ ID NO:4	biohr	31.55	
SEQ ID NO:4+biotin	biohr	40.51	7.5
SEQ ID NO:4+fluorescein	biohr	46.81	1.6
SEQ ID NO:4+Lys-Trp-Lys	biohr	36.66	6.1,
SEQ ID NO:4+o-phenanthroline	biohr	36.66	19.1
10 SEQ ID NO:4+pyrene	biohr	54.29	23.2
SEQ ID NO:4+TEA	biohr	31.81	5.9
SEQ ID NO:3	anal.	17.59	
SEQ ID NO:4	anal.	17.30	
SEQ ID NO:4+PEHA	anal.	17.34	1.8
15 SEQ ID NO:4+Lys-Tyr-Lys	anal.	17.37	8.6
SEQ ID NO:4+PEG	anal.	36.00	9.3

The "biohr" HPLC gradient was as follows: Solvent A: 50 mM TEAA, pH=7.4; solvent B: CH₃CN; 0-10 mins., 95% A, 5% B; linear increase to 15% B in the next fifty mins. HPLC column: Water's Delta-Pak C-18, reverse phase. The "anal." HPLC gradient was as follows: Solvent A: 50 mM TEAA, pH=7.4; Solvent B: CH₃CN; 0-10 mins., 95% A, 5% B; linear increase to 60% B in the next fifty mins. HPLC column: Water's Delta-Pak C-18, reverse phase.

25 **B. Oligonucleotide Having SEQ ID NO:16.**

The oligonucleotide having sequence GANCT (Seq. ID No. 16) (40 ODS) was treated in 100 μL of 1M NaOAC solution followed by 10 mg of 5-amino-O-phenanthroline dissolved in 200 μL of 1M NaOAC. After about 15 mts., 100 μL of a NaCNBH₃ solution (0.2M NaBH₃CN in 0.25M NaOAC) was added to the reaction and allowed to stand overnight. The conjugate was purified by size exclusion and reverse-phase HPLC.

- 54 -

TABLE XXIX

HPLC Data	HPLC program	Retention Time (min.)
SEQ ID NO: 15 I-6839	biohr	32.39
5 SEQ ID NO: 16 I-6839-D	biohr	27.25
I-6839-OP	biohr	47.87

The NMR spectra show the formation of the abasic site and the conjugate between phenanthroline ligand and the pentamer oligonucleotide. The final product is a homogeneous single product with no DNA fragmentation or 1,4-addition products evidenced.

¹H NMR analysis showed the following peaks: In the case of 6839 and 6839D, peaks between 7.4 and 8.4; whereas in 6839-OP peaks between 7.0 and 8.8 (protons from O-phenanthroline). In other words, 6839-OP conjugate showed the combination spectrum of Sequence 16 and O-phenanthroline. In ³¹P NMR dispersion of signals was higher for the conjugate (-0.7 to 0.4 ppm) than the Sequence 16.

EXAMPLE 14

To further derivatize the oligonucleotide-polyamine conjugate, imidazole-4-acetic acid is attached to the free amines made available by the polyamines attached in Example 4-A-4-b.

Imidazole-4-acetic acid is treated with 2,4-dinitrofluorobenzene. The product is treated with pentafluorophenol/DCC to give the active ester of imidazole-4-

- 55 -

acetic acid, which is also protected in the imidazole ring by a DNP group (Compound 13).

The oligonucleotide-polyamine conjugate is reacted with Compound 13 in 0.2M NaHCO₃ buffer/DMF. The product, 5 oligonucleotide-polyimidazole conjugate, then is treated with mercapto ethanol to remove the DNP group, and then is purified by size exclusion and HPLC methods.

- 56 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANTS: Manoharan, Muthiah
Phillip D. Cook
- (ii) TITLE OF INVENTION: NOVEL AMINES AND METHODS OF
MAKING AND USING THE SAME
- (iii) NUMBER OF SEQUENCES: 16
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Woodcock Washburn Kurtz Mackiewicz
and Norris
 - (B) STREET: One Liberty Place - 46th Floor
 - (C) CITY: Philadelphia
 - (D) STATE: PA
 - (E) COUNTRY: U.S.A.
 - (F) ZIP: 19103
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Gaumond, Rebecca R.
 - (B) REGISTRATION NUMBER: 35,152
 - (C) REFERENCE/DOCKET NUMBER: ISIS-1171
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 215-568-3100
 - (B) TELEFAX: 215-568-3439

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TGGGAGCCAT AGCGAGGCUCG

- 57 -

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 19
- (D) OTHER INFORMATION: /note= "abasic, aldehydic species"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TGGGAGCCAT AGCGAGGCN

19

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CGCAGUCAGC C

11

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6
- (D) OTHER INFORMATION: /note= "abasic residue"

- 58 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CGCAGNCAGC C

11

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 9
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 18
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GACAGAGGUA GGAGAAGUGA

20

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 9
- (D) OTHER INFORMATION: /note= "abasic residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 18
- (D) OTHER INFORMATION: /note= "abasic residue"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GACAGAGGNA GGAGAAGNGA

20

- 59 -

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TGGGAGCCAT AGCGAGGC

18

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 9
- (D) OTHER INFORMATION: /note=
"2'-O-aminopentoxo-2'-deoxyadenosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CTGTCTCCAT CCTCTTCACT

20

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CTGTCTCCAT CCTCTTCACT

20

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid

- 60 -

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 7
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 21
- (D) OTHER INFORMATION: /note= "2'-aminopropoxy cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGCGUCUCCA GGCGAUCUGA C

21

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 20
- (D) OTHER INFORMATION: /note= "2'-aminopropoxy cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TCTGAGTAGC AGAGGAGCTC

20

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 61 -

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 9
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 11
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 20
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 21
- (D) OTHER INFORMATION: /note= "2'-aminopropoxy cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GGAUGGCGUC UCCAGGCGAU C

21

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 9

- 62 -

(D) OTHER INFORMATION: /note= "2'deoxyuridine
residue"

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 11

(D) OTHER INFORMATION: /note= "2'deoxyuridine
residue"

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 20

(D) OTHER INFORMATION: /note= "2'deoxyuridine
residue"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GGAUGGCGUC UCCAGGCGAU C

21

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 18

(D) OTHER INFORMATION: /note= "2'deoxyuridine
residue"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TGGGAGCCAT AGCGAGGUCT

19

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 3

(D) OTHER INFORMATION: /note= "2'deoxyuridine
residue"

- 63 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GAUCT

5

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 3
- (D) OTHER INFORMATION: /note= "abasic residue"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:


GANCT

5

- 64 -

WHAT IS CLAIMED IS:

1. A compound having the structure:



wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic amine-containing species, and X is H, O- R_{11} , S- R_{11} , F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkyaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide.

2. The compound of claim 1 wherein the amine-containing species comprises at least one nitrogen atom having a free electron pair.


3. The compound of claim 1 wherein the R_1 and R_2 are each oligonucleotides.

- 65 -

4. The compound of claim 1 further including one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins or cross-linking agents attached to
5 at least one of the nitrogen atoms of said amine-containing species.

5. The compound of claim 1 in a pharmaceutically acceptable carrier.

6. A compound having the structure:



10 wherein R₄ is an oligonucleotide and M is a pendent group having an amine-containing species attached thereto.

7. The compound of claim 6 wherein M is R₇S⁻ or R₇NH⁺ wherein R₇ is an amine-containing species.


8. The compound of claim 7 wherein the amine-
15 containing species comprises at least one nitrogen atom having a free electron pair.

9. The compound of claim 6 further including one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides,
20 crown amines, porphyrins, or cross-linking agents attached to at least one of the nitrogen atoms of said amine-containing species.

10. The compound of claim 6 in a pharmaceutically acceptable carrier.

- 66 -

11. A compound having the structure:




wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic non-aromatic amine-containing species containing only non-amide nitrogen atoms, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle.

12. The compound of claim 11 wherein the amine-containing species comprises at least one nitrogen atom having a free electron pair.

13. The compound of claim 11 further including one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to at least one of the nitrogen atoms of said amine-containing species.

14. The compound of claim 11 in a pharmaceutically acceptable carrier.

15. A compound having the structure:



wherein B is a purine or pyrimidine heterocyclic base, R_8 and R_9 are independently H, PO_2^- , a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_8 and R_9 is a purine containing oligonucleotide; and at least one of R_8 , R_9 and A is a species comprising the formula L_1-L_2-

- 67 -

polyamine wherein L_1 is an amino linker and L_2 is a heterobifunctional linker;


wherein if R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- ; if R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH.

16. The compound of claim 15 wherein the amine-containing species comprises at least one nitrogen atom having a free electron pair.

17. The compound of claim 15 further including one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to at least one of the nitrogen atoms of said amine-containing species.

18. The compound of claim 15 in a pharmaceutically acceptable carrier.

19. A method of preparing a composition having the structure:




wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide, or an amine-containing species and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic non-aromatic amine-containing species, and X is H, $O-R_{11}$, $S-R_{11}$, F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter

- 68 -

molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkyaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide comprising the steps of:

providing a synthon having the structure:



wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide, or an amine-containing species and at least one of R_1 and R_2 is a purine containing oligonucleotide, and X is H, $O-R_{11}$, $S-R_{11}$, F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkyaryl or aralkyl, a ^{14}C containing C_7 to C_{14}

- 69 -

alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide;

and reacting said synthon with $-NH(R_3)$, wherein R_3 is a linear or cyclic non-aromatic amine-containing species, under reducing conditions to yield said compound.

20. The method of claim 19 wherein the amine-containing species has the formula $H_2N[(CH_2)_nNH]_m-$ wherein n is an integer between 2 and 8 and m is an integer between 1 and 10.


21. The method of claim 19 wherein the synthon is prepared by reacting an oligonucleotide with an appropriate DNA glycosylase.

22. The method of claim 19 wherein R_3 is a crown amine.

23. The method of claim 19 further comprising complexing said compound with a metal ion.

24. A novel compound prepared in accordance with claim 19.


25. A method of preparing a compound having the structure:



- 70 -

wherein R_4 is an oligonucleotide and M is a pendent group having an amine-containing species attached thereto comprising the steps of:

providing a compound having the structure:



5 wherein R_4 is an oligonucleotide, R_{12} is an oligonucleotide and B is urea or a heterocyclic base having a corresponding glycosylase;

reacting the compound with an endonuclease to generate a conjugated α, β -unsaturated system in the sugar
10 residue of the 3' terminal nucleotide;

reacting the compound with a pendent group containing a nucleophilic functionality thereon;

reacting the compound with a reducing agent to stabilize the compound; and

15 reacting the compound with an amine-containing species to add a pendent amine-containing species to said pendent group.

26. The method of claim 25 wherein said amine-containing species is added to said pendent group via an
20 alkylation reaction.

27. The method of claim 25 wherein said pendent group is a bifunctional linker.

28. The method of claim 25 wherein the nucleophile is $R_{10}S^-$ or $R_{10}NH^-$ wherein R_{10} is a said pendent group.


25 29. The method of claim 25 wherein the amine-containing species comprises at least one nitrogen group have a free electron pair.

- 71 -

30. The method of claim 25 further including adding one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents to at least one of the nitrogen atoms of said amine-containing species.

31. The method of claim 25 wherein the endonuclease is selected from the group consisting of endonuclease III, T4 UV endonuclease and *M. luteus* UV endonuclease.

32. A method of preparing a compound having the structure:



wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic amine-containing species containing only non-amide nitrogen atoms, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle comprising the steps of:

reacting an oligonucleotide having a 3' ribofuranosyl sugar with an oxidizing agent to produce an activated dialdehyde-terminated oligonucleotide; and
reacting said activated oligonucleotide with a linear or cyclic amine-containing species under reducing conditions to yield said compound.

33. The method of claim 32 wherein the oxidizing agent is *m*-periodate.


34. The method of claim 32 wherein the amine-containing species has the formula $H_2N[(CH_2)_nNH]_m-$ wherein n is an integer between 2 and 8 and m is an integer between 1 and 10.

- 72 -

35. A novel compound prepared in accordance with claim 32.

36. A method for modulating the production of a protein by an organism comprising:

5 contacting an organism with a compound having the structure:



wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide or an amine-containing species and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is
 10 a linear or cyclic amine-containing species, and X is H, O- R_{11} , S- R_{11} , F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the
 15 pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower
 20 alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkyaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter
 25 molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide.

- 73 -


37. The method of claim 36 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.

38. The method of claim 36 wherein said compound is hybridizable with an RNA coding for said protein.

39. The method of claim 36 wherein R_1 and R_2 , taken together range from about 8 to about 50 nucleotide bases in length.

40. The method of claim 36 wherein R_1 and R_2 taken together range from about 12 to about 20 nucleotide bases in length.

41. A method of treating an animal having a disease characterized by undesired production of protein comprising: contacting an animal with a compound having the structure:



wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic amine-containing species, and X is H, O- R_{11} , S- R_{11} , F, Cl, Br, CN, CF_3 , OCF_3 , OCN, SOCH_3 , SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an

- 74 -

oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide.

42. The method of claim 41 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.

43. The method of claim 41 wherein said compound is hybridizable with an RNA coding for said protein.


44. The method of claim 41 wherein R_1 and R_2 , taken together range from about 8 to about 50 nucleotide bases in length.

45. The method of claim 41 wherein R_1 and R_2 taken together range from about 12 to about 20 nucleotide bases in length.

46. A method for detecting the presence or absence of an RNA in a biological sample suspected of containing said RNA comprising:

contacting a sample with a compound having the structure:

- 75 -



wherein R_1 and R_2 are independently H, a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic amine-containing species, and X is H, O-
 5 R_{11} , S- R_{11} , F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocylcoalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group
 10 for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or
 15 substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkyaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the
 20 pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide; said compound being specifically hybridizable with said RNA; and

detecting the presence or absence of hybridization
 25 of the compound to the sample wherein hybridization is indicative of the presence of RNA in the sample.

- 76 -


47. The method of claim 46 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.

48. The method of claim 46 wherein R_1 and R_2 , taken together range from about 8 to about 50 nucleotide bases in length.

49. The method of claim 46 wherein R_1 and R_2 taken together range from about 12 to about 20 nucleotide bases in length.

50. A method for modulating the production of a protein by an organism comprising:

contacting an organism with a compound having the structure:



wherein R_4 is an oligonucleotide and M is a pendent group having an amine-containing species attached thereto.

51. The method of claim 50 wherein the compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.


52. The method of claim 50 wherein said compound is hybridizable with an RNA coding for said protein.

- 77 -

53. The method of claim 50 wherein R_4 ranges from about 8 to about 50 nucleotide bases in length.

54. The method of claim 50 wherein R_4 ranges from about 12 to about 20 nucleotide bases in length.

5 55. A method of treating an animal having a disease characterized by undesired production of protein comprising:
contacting an animal with a compound having the structure:



wherein R_4 is an oligonucleotide and M is a pendent group
10 having an amine-containing species attached thereto, in a pharmaceutically acceptable carrier.

56. The method of claim 55 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules,
15 steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.

57. The method of claim 55 wherein said compound is hybridizable with an RNA coding for said protein.


58. The method of claim 55 wherein R_4 ranges from
20 about 8 to about 50 nucleotide bases in length.

59. The method of claim 55 wherein R_4 ranges from about 12 to about 20 nucleotide bases in length.

- 78 -

60. A method for detecting the presence or absence of an RNA in a biological sample suspected of containing said RNA comprising:

contacting a sample with a compound having the
5 structure:



wherein R₄ is an oligonucleotide and M is a pendent group having an amine-containing species attached thereto; said compound being specifically hybridizable with said RNA; and

detecting the presence or absence of hybridization
10 of the compound to the sample wherein hybridization is indicative of the presence of RNA in the sample.

61. The method of claim 60 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking
15 agents attached to said amine-containing species.


62. The method of claim 60 wherein R₄ ranges from about 8 to about 50 nucleotide bases in length.

63. The method of claim 60 wherein R₄ ranges from
20 about 12 to about 20 nucleotide bases in length.

64. A method for modulating the production of a protein by an organism comprising:


contacting an organism with a compound having the structure:

- 79 -



wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic amine-containing species containing only non-amide nitrogen atoms, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle.

- 5 65. The method of claim 64 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.
- 10 66. The method of claim 64 wherein said compound is hybridizable with an RNA coding for said protein.
67. The method of claim 64 wherein R_4 and R_6 taken together range from 8 to about 50 nucleotide bases in length.
68. The method of claim 64 wherein R_4 and R_6 taken
15 together range from 12 to about 20 nucleotide bases in length.
69. A method of treating an animal having a disease characterized by undesired production of protein comprising contacting the animal with a compound having the structure:



- wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic
20 amine-containing species containing only non-amide nitrogen atoms, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle, in a pharmaceutically acceptable carrier.

- 80 -

70. The method of claim 69 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.


71. The method of claim 69 wherein said compound is hybridizable with an RNA coding for said protein.

72. The method of claim 69 wherein R_4 and R_6 taken together range from 8 to about 50 nucleotide bases in length.

73. The method of claim 69 wherein R_4 and R_6 taken together range from 12 to about 20 nucleotide bases in length.

74. A method for detecting the presence or absence of an RNA in a biological sample suspected of containing said RNA comprising:

contacting the sample with a compound having the structure:



wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic amine-containing species containing only non-amide nitrogen atoms, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle; said compound being specifically hybridizable with RNA; and

detecting hybridization of the compound to the sample where the presence or absence of hybridization is indicative of the presence of said RNA in the sample.


- 81 -

75. The method of claim 74 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.

76. The method of claim 74 wherein R_4 and R_6 taken together range from 8 to about 50 nucleotide bases in length.

77. The method of claim 74 wherein R_4 and R_6 taken together range from 12 to about 20 nucleotide bases in length.

78. A method for modulating the production of a protein by an organism comprising:
contacting an organism with a compound having the structure:



wherein B is a purine or pyrimidine heterocycle, R_8 and R_9 are independently H, PO_2^- , a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_8 and R_9 is a purine containing oligonucleotide, and at least one of R_8 , R_9 and A is a species comprising the formula L_1 - L_2 -polyamine wherein L_1 is an amino linker and L_2 is a heterobifunctional linker;

wherein if R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- ; if R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH.

- 82 -


79. The method of claim 78 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.

80. The method of claim 78 wherein said compound is hybridizable with an RNA coding for said protein.

81. The method of claim 78 wherein R_8 and R_9 taken together range from about 8 to about 50 nucleotide bases in length.

82. The method of claim 79 wherein R_8 and R_9 taken together range from about 12 to about 20 nucleotide bases in length.

83. A method of treating an animal having a disease characterized by undesired production of protein comprising: contacting an animal with a compound having the structure:



wherein B is a purine or pyrimidine heterocycle, R_8 and R_9 are independently H, PO_2^- , a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_8 and R_9 is a purine containing oligonucleotide, and at least one of R_8 , R_9 and A is a species comprising the formula L_1 - L_2 -polyamine wherein L_1 is an amino linker and L_2 is a heterobifunctional linker, in a pharmaceutically acceptable carrier; and wherein if R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- ; if R_9 is not a purine containing oligonucleotide or

- 83 -

polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH.

84. The method of claim 83 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.


85. The method of claim 83 wherein said compound is hybridizable with an RNA coding for said protein.

86. The method of claim 83 wherein R_8 and R_9 taken together range from about 8 to about 50 nucleotide bases in length.

87. The method of claim 83 wherein R_8 and R_9 taken together range from about 12 to about 20 nucleotide bases in length.

88. A method for detecting the presence or absence of an RNA in a biological sample suspected of containing said RNA comprising:

contacting a sample with a compound having the structure:



wherein B is a purine or pyrimidine heterocycle, R_8 and R_9 are independently H, PO_2^- , a nucleotide, an oligonucleotide or an amine-containing species, and at least one of R_8 and R_9 is a purine containing oligonucleotide, and at least one of R_8 , R_9 and A is a species comprising the formula $\text{L}_1\text{-L}_2\text{-polyamine}$

- 84 -

wherein L_1 is an amino linker and L_2 is a heterobifunctional linker; and

wherein if R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- ; if R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH; said compound being specifically hybridizable with said RNA; and

detecting the presence or absence of hybridization of the compound to the sample wherein hybridization is indicative of the presence of RNA in the sample.

89. The method of claim 88 wherein said compound further includes one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to said amine-containing species.

90. The method of claim 88 wherein R_8 and R_9 taken together range from about 8 to about 50 nucleotide bases in length.

91. The method of claim 88 wherein R_8 and R_9 taken together range from about 12 to about 20 nucleotide bases in length.

CONJUGATION CHEMISTRY USING URACIL-DNA GLYCOSYLASE




FIGURE 1

2/3




FIGURE 2

3'- OLIGO POLYAMINE CONJUGATION




FIGURE 3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/08367

A. CLASSIFICATION OF SUBJECT MATTER IPC(5) : C07H 21/04; C12Q 1/68; A61K 48/00 US CL : 435/6; 536/23.1, 24.3, 24.5; 514/44 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6; 536/23.1, 24.3, 24.5; 514/44 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Extra Sheet.		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Journal of Biochemistry, Volume 101, issued 1987, H. Sawai et al, "Synthesis and Biological Activities of β -Alanyltyrosine Derivative of 2',5'-Oligoadenylate, and Its Use in Radiobinding Assay for 2',5'-Oligoadenylate", pages 339-346, especially paragraph bridging pages 340-341 and Figure 1.	11-14, 32-35, 64-77
Y	Journal of Biochemistry, Volume 98, issued 1985, H. Sawai et al, "Sensitive Radioimmuno Assay for 2',5'-Oligoadenylates Using a Novel ¹²⁵ I-Labeled Derivative of 2',5'-Triadenylate 5'-Triphosphate", pages 999-1005, especially page 1000, third full paragraph and Figure 1.	11-14, 32-35, 64-77
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* "A" "E" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be part of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to underpin the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combinations being obvious to a person skilled in the art "Z" document member of the same patent family
Date of the actual completion of the international search 10 DECEMBER 1993		Date of mailing of the international search report 20 DEC 1993
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. NOT APPLICABLE		Authorized officer DAVID SCHREIBER <i>DP. Schreiber</i> Telephone No. (703) 308-0196

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Methods in Enzymology, Volume 18, Pt. B, issued 1971, W. Föry et al, "Chemical Synthesis of Flavin Coenzymes", pages 458-464, especially Figures VII and VIII.	1-10, 36-63
Y	Pharmaceutical Research, Volume 5. No. 9, issued 1988, G. Zon, "Oligonucleotide Analogues as Potential Chemotherapeutic Agents", pages 539-549, entire document.	36-91
Y	Bioconjugate Chemistry, Volume 1, Number 3, issued May/June 1990, J. Goodchild, "Conjugates of Oligonucleotides and Modified Oligonucleotides: A Review of their Synthesis and Properties", pages 165-187, entire document.	11-14, 36-91
Y	US, A, 5,034,506 (Summerton et al) 23 July 1991, claim 3.	11-14, 32-35, 64-77
Y	Mutation Research, Volume 236, issued 1990, P. W. Doetsch et al, "The enzymology of apurinic/apyrimidinic endonucleases", pages 173-201, especially Figure 3 and page 177, first full paragraph.	25-31
Y	Nucleosides & Nucleotides, Volume 10, Numbers 1-3, issued 1991, J.-J. Vasseur et al, "Derivatization of Oligonucleotides through Abasic Site Formation", pages 107-117, especially Figures 3 and 4.	25-31
Y	Proceedings of the National Academy of Sciences USA, Volume 86, issued September 1989, R. L. Letsinger et al, "Cholesteryl-conjugated oligonucleotides: Synthesis, properties, and activity as inhibitors of replication of human immunodeficiency virus in cell culture", pages 6553-6656, entire document.	4, 9, 13, 17, 30, 37, 42, 47, 51, 56, 61, 65, 70, 75, 79, 84, 89
Y	Journal of the American Chemical Society, Volume 110, issued 1988, M. Manoharan et al, "Mechanism of UV Endonuclease V Cleavage of Abasic Sites in DNA Determined by ¹³ C Labeling", pages 2690-2691, especially Scheme I.	25-31

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/08367

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
(Form PCT/ISA/206 Previously Mailed.)
Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☒ No protest accompanied the payment of additional search fees.

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

CAS ONLINE, MEDLINE, APS, BIOSIS, WPI,

search terms: structure search, dialdehyde, periodate, amine, amino, polyamine, morpholino, nucleic, DNA, RNA, ribonucleic, deoxyribonucleic, heterobifunctional, crown amine, steroids, porphyrins, nucleotide, nucleoside, oligonucleotide, detection, productin, modulate, protein

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

- I. Claims 1-5, 19-24, and 36-45, drawn to a compound having the structure shown in independent claim 1, a method for making this compound, and a method for modulating the production of a protein by an organism using this compound, classified in Class 536, subclass 23.1, for example;
- II. Claims 46-49, drawn to a method for detecting the presence or absence of an RNA in a biological sample using a compound having the structure shown in independent claim 1, classified in Class 435, subclass 6;
- III. Claims 6-10, 25-31, and 50-59, drawn to a compound having the structure shown in independent claim 6, a method for making this compound, and a method for modulating the production of a protein by an organism using this compound, classified in Class 536, subclass 23.1, for example;
- IV. Claims 60-63, drawn to a method for detecting the presence or absence of an RNA in a biological sample using a compound having the structure shown in independent claim 6, classified in Class 435, subclass 6;
- V. Claims 11-14, 32-35, and 64-73, drawn to a compound having the structure shown in independent claim 11, a method for making this compound, and a method for modulating the production of a protein by an organism using this compound, classified in Class 536, subclass 23.1, for example;
- VI. Claims 74-77, drawn to a method for detecting the presence or absence of an RNA in a biological sample using a compound having the structure shown in independent claim 11, classified in Class 435, subclass 6;
- VII. Claims 15-18 and 78-87, drawn to a compound having the structure shown in independent claim 15 and a method for modulating the production of a protein by an organism using this compound, classified in Class 536, subclass 23.1, for example;
- VIII. Claims 88-91, drawn to a method for detecting the presence or absence of an RNA in a biological sample using a compound having the structure shown in independent claim 15, classified in Class 435, subclass 6